


1A P20 Rec'd 24 MAR 2006

HIV REPLICATION INHIBITING PURINE DERIVATIVES


- The present invention concerns purine derivatives having HIV (Human Immunodeficiency Virus) replication inhibiting properties. The invention further relates to methods for their preparation and pharmaceutical compositions comprising them. The invention also relates to the use of said compounds for the manufacture of a medicament for the prevention or the treatment of HIV infection.
- WO 98/05335 describes 2,6,9-trisubstituted purine derivatives as inhibitors of cyclin dependent kinase 2 and I κ B- α .
WO 01/09134 describes purine derivatives as inhibitors of tyrosine protein kinase SYK.
- The compounds of the invention differ from the prior art compounds in structure, pharmacological activity, pharmacological potency.
- Unexpectedly, it has been found that the compounds of the invention have an improved ability to inhibit the replication of HIV, including an improved ability to inhibit the replication of mutant strains, i.e. strains which have become resistant to art-known drug(s) (drug or multidrug resistant HIV strains).

The present invention concerns the use of a compound of formula (I) for the manufacture of a medicament for the prevention or the treatment of HIV infection wherein the compound of formula (I) is a compound of formula



a *N*-oxide, a pharmaceutically acceptable addition salt, a quaternary amine or a stereochemically isomeric form thereof, wherein

A and B each represents a radical of formula



wherein

- ring E represents phenyl, pyridyl, pyridazinyl, pyrimidinyl or pyrazinyl;
ring F represents phenyl, pyridyl, pyridazinyl, pyrimidinyl or pyrazinyl;

-2-

- R^1 represents hydrogen; aryl; formyl; C_{1-6} alkylcarbonyl; C_{1-6} alkyloxycarbonyl; C_{1-6} alkyl optionally substituted with formyl, C_{1-6} alkylcarbonyl, C_{1-6} alkyloxycarbonyl, C_{1-6} alkylcarbonyloxy; or C_{1-6} alkyloxy C_{1-6} alkylcarbonyl substituted with C_{1-6} alkyloxycarbonyl;
- 5 R^2 represents cyano; aminocarbonyl; mono- or di(C_{1-4} alkyl)aminocarbonyl; C_{1-6} alkyl optionally substituted with cyano, aminocarbonyl or mono- or di(C_{1-4} alkyl)aminocarbonyl; C_{2-6} alkenyl substituted with cyano, aminocarbonyl or mono- or di(C_{1-4} alkyl)aminocarbonyl; or C_{2-6} alkynyl substituted with cyano, aminocarbonyl or mono- or di(C_{1-4} alkyl)aminocarbonyl;
- 10 X_1 represents $-NR^5-$; $-NH-NH-$; $-N=N-$; $-O-$; $-C(=O)-$; $-C_{1-4}$ alkanediyl-; $-CHOH-$; $-S-$; $-S(=O)_p-$; $-X_2-C_{1-4}$ alkanediyl-; $-C_{1-4}$ alkanediyl- X_2- ; or $-C_{1-4}$ alkanediyl- X_2-C_{1-4} alkanediyl-;
- X_2 represents $-NR^5-$; $-NH-NH-$; $-N=N-$; $-O-$; $-C(=O)-$; $-CHOH-$; $-S-$; or $-S(=O)_p-$;
- m represents an integer of value 1, 2, 3 or 4;
- 15 R^3 represents cyano; aminocarbonyl; amino; halo; NHR^{13} ; $NR^{13}R^{14}$; $-C(=O)-NHR^{13}$; $-C(=O)-NR^{13}R^{14}$; $-C(=O)-R^{15}$; $-CH=N-NH-C(=O)-R^{16}$; C_{1-6} alkyl optionally substituted with one or more substituents each independently selected from R^{3a} ; C_{1-6} alkyloxy optionally substituted with one or more substituents each independently selected from R^{3a} ; C_{1-6} alkyloxy C_{1-6} alkyl optionally substituted with one or more substituents each independently selected from R^{3a} ; C_{2-6} alkenyl optionally substituted with one or more substituents each independently selected from R^{3a} ; C_{2-6} alkynyl optionally substituted with one or more substituents each independently selected from R^{3a} ; $-C(=N-O-R^8)-C_{1-4}$ alkyl; R^7 or $-X_3-R^7$;
- 20 R^{3a} represents halo, cyano, hydroxy, NR^9R^{10} , $-C(=O)-NR^9R^{10}$, $-C(=O)-C_{1-6}$ alkyl, $-C(=O)-O-C_{1-6}$ alkyl, $-C(=O)-polyhaloC_{1-6}$ alkyl, $-C(=O)-O-polyhaloC_{1-6}$ alkyl or R^7 ;
- 25 X_3 represents $-NR^5-$; $-NH-NH-$; $-N=N-$; $-O-$; $-C(=O)-$; $-S-$; $-S(=O)_p-$; $-X_{4a}-C_{1-4}$ alkanediyl-; $-C_{1-4}$ alkanediyl- $X_{4b}-$; $-C_{1-4}$ alkanediyl- $X_{4a}-C_{1-4}$ alkanediyl-; or $-C(=N-OR^8)-C_{1-4}$ alkanediyl-;
- X_{4a} represents $-NR^5-$; $-NH-NH-$; $-N=N-$; $-C(=O)-$; $-S-$; or $-S(=O)_p-$;
- 30 X_{4b} represents $-NH-NH-$; $-N=N-$; $-O-$; $-C(=O)-$; $-S-$; or $-S(=O)_p-$;
- each R^4 independently represents hydroxy; halo; C_{1-6} alkyl optionally substituted with one or more substituents each independently selected from R^{4a} ; C_{2-6} alkenyl optionally substituted with one or more substituents each independently selected from R^{4a} ; C_{2-6} alkynyl optionally substituted with one or more substituents each independently selected from R^{4a} ; C_{3-7} cycloalkyl; C_{1-6} alkyloxy; C_{1-6} alkyloxycarbonyl; C_{1-6} alkylcarbonyloxy; carboxyl; formyl; cyano; nitro; amino; mono- or
- 35 di(C_{1-6} alkyl)amino; polyhalo C_{1-6} alkyl; polyhalo C_{1-6} alkyloxy; polyhalo C_{1-6} alkylthio;

-3-

$-S(=O)_pR^6$; $-NH-S(=O)_pR^6$; $-C(=O)R^6$; $-NHC(=O)H$; $-C(=O)NHNH_2$; $NHC(=O)R^6$; $C(=NH)R^6$; or R^7 ;

R^{4a} represents halo, cyano, NR^9R^{10} , hydroxy or $-C(=O)R^6$;

R^5 represents hydrogen; aryl; formyl; C_{1-6} alkylcarbonyl; C_{1-6} alkyloxycarbonyl;

5 C_{1-6} alkyl optionally substituted with formyl, C_{1-6} alkylcarbonyl, C_{1-6} alkyloxycarbonyl or C_{1-6} alkylcarbonyloxy; or C_{1-6} alkyloxy C_{1-6} alkylcarbonyl substituted with C_{1-6} alkyloxycarbonyl;

R^6 represents C_{1-6} alkyl, amino, mono- or di(C_{1-4} alkyl)amino or polyhalo C_{1-4} alkyl;

R^7 represents a monocyclic, bicyclic or tricyclic saturated carbocycle; a monocyclic,
10 bicyclic or tricyclic partially saturated carbocycle; a monocyclic, bicyclic or tricyclic aromatic carbocycle; a monocyclic, bicyclic or tricyclic saturated heterocycle; a monocyclic, bicyclic or tricyclic partially saturated heterocycle; or a monocyclic, bicyclic or tricyclic aromatic heterocycle; wherein each of said carbocyclic or heterocyclic ring systems may, whenever possible, optionally be substituted with
15 one, two, three, four or five substituents each independently selected from halo, hydroxy, mercapto, C_{1-6} alkyl, hydroxy C_{1-6} alkyl, amino C_{1-6} alkyl, mono or di(C_{1-6} alkyl)amino C_{1-6} alkyl, formyl, C_{1-6} alkylcarbonyl, C_{3-7} cycloalkyl, C_{1-6} alkyloxy, C_{1-6} alkyloxycarbonyl, C_{1-6} alkylthio, cyano, nitro, polyhalo C_{1-6} alkyl, polyhalo C_{1-6} alkyloxy, aminocarbonyl, $-CH(=N-O-R^8)$, R^{7a} ,
20 $-X_3-R^{7a}$ or $R^{7a}-C_{1-4}$ alkanediyl-;

R^{7a} represents a monocyclic, bicyclic or tricyclic saturated carbocycle; a monocyclic, bicyclic or tricyclic partially saturated carbocycle; a monocyclic, bicyclic or tricyclic aromatic carbocycle; a monocyclic, bicyclic or tricyclic saturated heterocycle; a monocyclic, bicyclic or tricyclic partially saturated heterocycle; or a monocyclic,
25 bicyclic or tricyclic aromatic heterocycle; wherein each of said carbocyclic or heterocyclic ring systems may optionally be substituted with one, two, three, four or five substituents each independently selected from halo, hydroxy, mercapto, C_{1-6} alkyl, hydroxy C_{1-6} alkyl, amino C_{1-6} alkyl, mono or di(C_{1-6} alkyl)amino C_{1-6} alkyl, formyl, C_{1-6} alkylcarbonyl, C_{3-7} cycloalkyl, C_{1-6} alkyloxy, C_{1-6} alkyloxycarbonyl, C_{1-6} alkylthio, cyano, nitro, polyhalo C_{1-6} alkyl, polyhalo C_{1-6} alkyloxy, aminocarbonyl, $-CH(=N-O-R^8)$;

R^8 represents hydrogen, C_{1-4} alkyl optionally substituted with aryl, or aryl;

R^9 and R^{10} each independently represent hydrogen; hydroxy; C_{1-6} alkyl; C_{1-6} alkyloxy; C_{1-6} alkylcarbonyl; C_{1-6} alkyloxycarbonyl; amino; mono- or di(C_{1-6} alkyl)amino;
35 mono- or di(C_{1-6} alkyl)aminocarbonyl; $-CH(=NR^{11})$ or R^7 , wherein each of the aforementioned C_{1-6} alkyl groups may optionally and each individually be substituted with one or two substituents each independently selected from hydroxy,

-4-

C₁₋₆alkyloxy, hydroxyC₁₋₆alkyloxy, carboxyl, C₁₋₆alkyloxycarbonyl, cyano, amino, imino, mono- or di(C₁₋₄alkyl)amino, polyhaloC₁₋₄alkyl, polyhaloC₁₋₄alkyloxy, polyhaloC₁₋₄alkylthio, -S(=O)_pR⁶, -NH-S(=O)_pR⁶, -C(=O)R⁶, -NHC(=O)H, -C(=O)NHNH₂, -NHC(=O)R⁶, -C(=NH)R⁶, or R⁷; or

5 R⁹ and R¹⁰ may be taken together to form a bivalent or trivalent radical of formula

-CH₂-CH₂-CH₂-CH₂- (d-1);

-CH₂-CH₂-CH₂-CH₂-CH₂- (d-2);

-CH₂-CH₂-O-CH₂-CH₂- (d-3);

-CH₂-CH₂-S-CH₂-CH₂- (d-4);

10 -CH₂-CH₂-NR¹²-CH₂-CH₂- (d-5);

-CH₂-CH=CH-CH₂- (d-6); or

=CH-CH=CH-CH=CH- (d-7);

R¹¹ represents cyano; C₁₋₄alkyl optionally substituted with C₁₋₄alkyloxy, cyano, amino, mono- or di(C₁₋₄alkyl)amino or aminocarbonyl; C₁₋₄alkylcarbonyl;

15 C₁₋₄alkyloxycarbonyl; aminocarbonyl; mono- or di(C₁₋₄alkyl)aminocarbonyl;

R¹² represents hydrogen or C₁₋₄alkyl;

R¹³ and R¹⁴ each independently represent C₁₋₆alkyl optionally substituted with cyano, aminocarbonyl or mono- or di(C₁₋₄alkyl)aminocarbonyl; C₂₋₆alkenyl optionally substituted with cyano, aminocarbonyl or mono- or di(C₁₋₄alkyl)aminocarbonyl;

20 C₂₋₆alkynyl optionally substituted with cyano, aminocarbonyl or mono- or di(C₁₋₄alkyl)aminocarbonyl;

R¹⁵ represents C₁₋₆alkyl optionally substituted with cyano, aminocarbonyl or mono- or di(C₁₋₄alkyl)aminocarbonyl;

R¹⁶ represents C₁₋₆alkyl optionally substituted with cyano, aminocarbonyl or mono- or
25 di(C₁₋₄alkyl)aminocarbonyl; or R⁷;

-C-D- represents a bivalent radical of formula

-N=CH-NR¹⁷- (c-1); or

-NR¹⁷-CH=N- (c-2);

R¹⁷ represents hydrogen; C₁₋₆alkyl optionally substituted with hydroxy, cyano,
30 aminocarbonyl, mono- or di(C₁₋₄alkyl)aminocarbonyl, C₁₋₄alkyloxycarbonyl or aryl;
p represents an integer of value 1 or 2;

aryl represents phenyl or phenyl substituted with one, two, three, four or five substituents each independently selected from halo, hydroxy, mercapto, C₁₋₆alkyl, hydroxyC₁₋₆alkyl, aminoC₁₋₆alkyl, mono or di(C₁₋₆alkyl)aminoC₁₋₆alkyl,


35 C₁₋₆alkylcarbonyl, C₃₋₇cycloalkyl, C₁₋₆alkyloxy, C₁₋₆alkyloxycarbonyl, C₁₋₆alkylthio, cyano, nitro, polyhaloC₁₋₆alkyl, polyhaloC₁₋₆alkyloxy, aminocarbonyl, R⁷ or -X₃-R⁷;

-5-


provided that when A represents a radical of formula (a) then B represents a radical of formula (b) and when A represents a radical of formula (b) then B represents a radical of formula (a).

- 5 The present invention also relates to a compound of formula (I) as defined hereinabove provided that when R^2 represents aminocarbonyl or mono- or di(C_{1-6} alkyl)aminocarbonyl then R^3 represents cyano; $-C(=O)-R^{15}$; $-CH=N-NH-C(=O)-R^{16}$; C_{1-6} alkyl substituted with one or more substituents each independently selected from R^{3b} ; C_{1-6} alkyloxy substituted with one or more substituents each independently selected from R^{3a} ; C_{1-6} alkyloxy C_{1-6} alkyl optionally substituted with one or more substituents each independently selected from R^{3a} ; C_{2-6} alkenyl optionally substituted with one or more substituents each independently selected from R^{3a} ; C_{2-6} alkynyl optionally substituted with one or more substituents each independently selected from R^{3a} ; $-C(=N-O-R^8)-C_{1-4}$ alkyl; R^7 or $-X_3-R^7$; with R^{3b} representing cyano, hydroxy, NR^9R^{10} , $-C(=O)-NR^9R^{10}$, $-C(=O)-C_{1-6}$ alkyl, $-C(=O)-polyhaloC_{1-6}$ alkyl, $-C(=O)-O-polyhaloC_{1-6}$ alkyl or R^7 .

Thus, the present invention also relates to a compound of formula



- 20 a *N*-oxide, a pharmaceutically acceptable addition salt, a quaternary amine or a stereochemically isomeric form thereof, wherein A and B each represents a radical of formula



- ring E represents phenyl, pyridyl, pyridazinyl, pyrimidinyl or pyrazinyl;
25 ring F represents phenyl, pyridyl, pyridazinyl, pyrimidinyl or pyrazinyl;
 R^1 represents hydrogen; aryl; formyl; C_{1-6} alkylcarbonyl; C_{1-6} alkyloxycarbonyl; C_{1-6} alkyl optionally substituted with formyl, C_{1-6} alkylcarbonyl, C_{1-6} alkyloxycarbonyl, C_{1-6} alkylcarbonyloxy; or C_{1-6} alkyloxy C_{1-6} alkylcarbonyl substituted with C_{1-6} alkyloxycarbonyl;
30 R^2 represents cyano; aminocarbonyl; mono- or di(C_{1-4} alkyl)aminocarbonyl; C_{1-6} alkyl optionally substituted with cyano, aminocarbonyl or mono- or di(C_{1-4} alkyl)aminocarbonyl; C_{2-6} alkenyl substituted with cyano, aminocarbonyl or

-6-

- mono- or di(C₁₋₄alkyl)aminocarbonyl; or C₂₋₆alkynyl substituted with cyano, aminocarbonyl or mono- or di(C₁₋₄alkyl)aminocarbonyl;
- X₁ represents -NR⁵-; -NH-NH-; -N=N-; -O-; -C(=O)-; -C₁₋₄alkanediyl-; -CHOH-; -S-; -S(=O)_p-; -X₂-C₁₋₄alkanediyl-; -C₁₋₄alkanediyl-X₂-; or
- 5 -C₁₋₄alkanediyl-X₂-C₁₋₄alkanediyl-;
- X₂ represents -NR⁵-; -NH-NH-; -N=N-; -O-; -C(=O)-; -CHOH-; -S-; or -S(=O)_p-;
- m represents an integer of value 1, 2, 3 or 4;
- R³ represents cyano; aminocarbonyl; amino; halo; NHR¹³; NR¹³R¹⁴; -C(=O)-NHR¹³; -C(=O)-NR¹³R¹⁴; -C(=O)-R¹⁵; -CH=N-NH-C(=O)-R¹⁶; C₁₋₆alkyl optionally
- 10 substituted with one or more substituents each independently selected from R^{3a}; C₁₋₆alkyloxy optionally substituted with one or more substituents each independently selected from R^{3a}; C₁₋₆alkyloxyC₁₋₆alkyl optionally substituted with one or more substituents each independently selected from R^{3a}; C₂₋₆alkenyl optionally substituted with one or more substituents each independently selected from R^{3a}; C₂₋₆alkynyl
- 15 optionally substituted with one or more substituents each independently selected from R^{3a}; -C(=N-O-R⁸)-C₁₋₄alkyl; R⁷ or -X₃-R⁷;
- R^{3a} represents halo, cyano, hydroxy, NR⁹R¹⁰, -C(=O)-NR⁹R¹⁰, -C(=O)-C₁₋₆alkyl, -C(=O)-O-C₁₋₆alkyl, -C(=O)-polyhaloC₁₋₆alkyl, -C(=O)-O-polyhaloC₁₋₆alkyl or R⁷;
- X₃ represents -NR⁵-; -NH-NH-; -N=N-; -O-; -C(=O)-; -S-; -S(=O)_p-;
- 20 -X_{4a}-C₁₋₄alkanediyl-; -C₁₋₄alkanediyl-X_{4b}-; -C₁₋₄alkanediyl-X_{4a}-C₁₋₄alkanediyl-; or -C(=N-OR⁸)-C₁₋₄alkanediyl-;
- X_{4a} represents -NR⁵-; -NH-NH-; -N=N-; -C(=O)-; -S-; or -S(=O)_p-;
- X_{4b} represents -NH-NH-; -N=N-; -O-; -C(=O)-; -S-; or -S(=O)_p-;
- each R⁴ independently represents hydroxy; halo; C₁₋₆alkyl optionally substituted with
- 25 one or more substituents each independently selected from R^{4a}; C₂₋₆alkenyl optionally substituted with one or more substituents each independently selected from R^{4a}; C₂₋₆alkynyl optionally substituted with one or more substituents each independently selected from R^{4a}; C₃₋₇cycloalkyl; C₁₋₆alkyloxy; C₁₋₆alkyloxycarbonyl; C₁₋₆alkylcarbonyloxy; carboxyl; formyl; cyano; nitro; amino; mono- or
- 30 di(C₁₋₆alkyl)amino; polyhaloC₁₋₆alkyl; polyhaloC₁₋₆alkyloxy; polyhaloC₁₋₆alkylthio; -S(=O)_pR⁶; -NH-S(=O)_pR⁶; -C(=O)R⁶; -NHC(=O)H; -C(=O)NHNH₂; NHC(=O)R⁶; C(=NH)R⁶; or R⁷;
- R^{4a} represents halo, cyano, NR⁹R¹⁰, hydroxy or -C(=O)R⁶;
- R⁵ represents hydrogen; aryl; formyl; C₁₋₆alkylcarbonyl; C₁₋₆alkyloxycarbonyl;
- 35 C₁₋₆alkyl optionally substituted with formyl, C₁₋₆alkylcarbonyl, C₁₋₆alkyloxycarbonyl or C₁₋₆alkylcarbonyloxy; or C₁₋₆alkyloxyC₁₋₆alkylcarbonyl substituted with C₁₋₆alkyloxycarbonyl;

-7-

R⁶ represents C₁₋₆alkyl, amino, mono- or di(C₁₋₄alkyl)amino or polyhaloC₁₋₄alkyl;



R⁷ represents a monocyclic, bicyclic or tricyclic saturated carbocycle; a monocyclic, bicyclic or tricyclic partially saturated carbocycle; a monocyclic, bicyclic or tricyclic aromatic carbocycle; a monocyclic, bicyclic or tricyclic saturated heterocycle; a monocyclic, bicyclic or tricyclic partially saturated heterocycle; or a monocyclic, bicyclic or tricyclic aromatic heterocycle; wherein each of said carbocyclic or heterocyclic ring systems may, whenever possible, optionally be substituted with one, two, three, four or five substituents each independently selected from halo, hydroxy, mercapto, C₁₋₆alkyl, hydroxyC₁₋₆alkyl, aminoC₁₋₆alkyl, mono or di(C₁₋₆alkyl)aminoC₁₋₆alkyl, formyl, C₁₋₆alkylcarbonyl, C₃₋₇cycloalkyl, C₁₋₆alkyloxy, C₁₋₆alkyloxycarbonyl, C₁₋₆alkylthio, cyano, nitro, polyhaloC₁₋₆alkyl, polyhaloC₁₋₆alkyloxy, aminocarbonyl, -CH(=N-O-R⁸), R^{7a}, -X₃-R^{7a} or R^{7a}-C₁₋₄alkanediyl-;

R^{7a} represents a monocyclic, bicyclic or tricyclic saturated carbocycle; a monocyclic, bicyclic or tricyclic partially saturated carbocycle; a monocyclic, bicyclic or tricyclic aromatic carbocycle; a monocyclic, bicyclic or tricyclic saturated heterocycle; a monocyclic, bicyclic or tricyclic partially saturated heterocycle; or a monocyclic, bicyclic or tricyclic aromatic heterocycle; wherein each of said carbocyclic or heterocyclic ring systems may optionally be substituted with one, two, three, four or five substituents each independently selected from halo, hydroxy, mercapto, C₁₋₆alkyl, hydroxyC₁₋₆alkyl, aminoC₁₋₆alkyl, mono or di(C₁₋₆alkyl)aminoC₁₋₆alkyl, formyl, C₁₋₆alkylcarbonyl, C₃₋₇cycloalkyl, C₁₋₆alkyloxy, C₁₋₆alkyloxycarbonyl, C₁₋₆alkylthio, cyano, nitro, polyhaloC₁₋₆alkyl, polyhaloC₁₋₆alkyloxy, aminocarbonyl, -CH(=N-O-R⁸);

R⁸ represents hydrogen, C₁₋₄alkyl optionally substituted with aryl, or aryl;

R⁹ and R¹⁰ each independently represent hydrogen; hydroxy; C₁₋₆alkyl; C₁₋₆alkyloxy; C₁₋₆alkylcarbonyl; C₁₋₆alkyloxycarbonyl; amino; mono- or di(C₁₋₆alkyl)amino; mono- or di(C₁₋₆alkyl)aminocarbonyl; -CH(=NR¹¹) or R⁷, wherein each of the aforementioned C₁₋₆alkyl groups may optionally and each individually be substituted with one or two substituents each independently selected from hydroxy, C₁₋₆alkyloxy, hydroxyC₁₋₆alkyloxy, carboxyl, C₁₋₆alkyloxycarbonyl, cyano, amino, imino, mono- or di(C₁₋₄alkyl)amino, polyhaloC₁₋₄alkyl, polyhaloC₁₋₄alkyloxy, polyhaloC₁₋₄alkylthio, -S(=O)_pR⁶, -NH-S(=O)_pR⁶, -C(=O)R⁶, -NHC(=O)H, -C(=O)NHNH₂, -NHC(=O)R⁶, -C(=NH)R⁶, or R⁷; or

R⁹ and R¹⁰ may be taken together to form a bivalent or trivalent radical of formula



-8-

- CH₂-CH₂-O-CH₂-CH₂- (d-3);
-CH₂-CH₂-S-CH₂-CH₂- (d-4);
-CH₂-CH₂-NR¹²-CH₂-CH₂- (d-5);
-CH₂-CH=CH-CH₂- (d-6); or
5 =CH-CH=CH-CH=CH- (d-7);

R¹¹ represents cyano; C₁₋₄alkyl optionally substituted with C₁₋₄alkyloxy, cyano, amino, mono- or di(C₁₋₄alkyl)amino or aminocarbonyl; C₁₋₄alkylcarbonyl;

C₁₋₄alkyloxycarbonyl; aminocarbonyl; mono- or di(C₁₋₄alkyl)aminocarbonyl;

R¹² represents hydrogen or C₁₋₄alkyl;

- 10 R¹³ and R¹⁴ each independently represent C₁₋₆alkyl optionally substituted with cyano, aminocarbonyl or mono- or di(C₁₋₄alkyl)aminocarbonyl; C₂₋₆alkenyl optionally substituted with cyano, aminocarbonyl or mono- or di(C₁₋₄alkyl)aminocarbonyl; C₂₋₆alkynyl optionally substituted with cyano, aminocarbonyl or mono- or di(C₁₋₄alkyl)aminocarbonyl;

- 15 R¹⁵ represents C₁₋₆alkyl optionally substituted with cyano, aminocarbonyl or mono- or di(C₁₋₄alkyl)aminocarbonyl;

R¹⁶ represents C₁₋₆alkyl optionally substituted with cyano, aminocarbonyl or mono- or di(C₁₋₄alkyl)aminocarbonyl; or R⁷;

-C-D- represents a bivalent radical of formula

- 20 -N=CH-NR¹⁷- (c-1); or
 -NR¹⁷-CH=N- (c-2);

R¹⁷ represents hydrogen; C₁₋₆alkyl optionally substituted with hydroxy, cyano, aminocarbonyl, mono- or di(C₁₋₄alkyl)aminocarbonyl, C₁₋₄alkyloxycarbonyl or aryl; p represents an integer of value 1 or 2;

- 25 aryl represents phenyl or phenyl substituted with one, two, three, four or five substituents each independently selected from halo, hydroxy, mercapto, C₁₋₆alkyl, hydroxyC₁₋₆alkyl, aminoC₁₋₆alkyl, mono or di(C₁₋₆alkyl)aminoC₁₋₆alkyl, C₁₋₆alkylcarbonyl, C₃₋₇cycloalkyl, C₁₋₆alkyloxy, C₁₋₆alkyloxycarbonyl, C₁₋₆alkylthio, cyano, nitro, polyhaloC₁₋₆alkyl, polyhaloC₁₋₆alkyloxy,
30 aminocarbonyl, R⁷ or -X₃-R⁷;

provided that when A represents a radical of formula (a) then B represents a radical of formula (b) and when A represents a radical of formula (b) then B represents a radical of formula (a); and

provided that when R² represents aminocarbonyl or mono- or

- 35 di(C₁₋₄alkyl)aminocarbonyl then R³ represents cyano; -C(=O)-R¹⁵;
-CH=N-NH-C(=O)-R¹⁶; C₁₋₆alkyl substituted with one or more substituents each independently selected from R^{3b}; C₁₋₆alkyloxy substituted with one or more substituents

-9-

each independently selected from R^{3a} ; C_{1-6} alkyloxy C_{1-6} alkyl optionally substituted with one or more substituents each independently selected from R^{3a} ; C_{2-6} alkenyl optionally substituted with one or more substituents each independently selected from R^{3a} ; C_{2-6} alkynyl optionally substituted with one or more substituents each independently selected from R^{3a} ; $-C(=N-O-R^8)-C_{1-4}$ alkyl; R^7 or $-X_3-R^7$; with R^{3b} representing cyano, hydroxy, NR^9R^{10} , $-C(=O)-NR^9R^{10}$, $-C(=O)-C_{1-6}$ alkyl, $-C(=O)-polyhaloC_{1-6}$ alkyl, $-C(=O)-O-polyhaloC_{1-6}$ alkyl or R^7 .

As used hereinbefore or hereinafter C_{1-4} alkyl as a group or part of a group defines straight or branched chain saturated hydrocarbon radicals having from 1 to 4 carbon atoms such as methyl, ethyl, propyl, 1-methylethyl, butyl; C_{1-6} alkyl as a group or part of a group defines straight or branched chain saturated hydrocarbon radicals having from 1 to 6 carbon atoms such as the group defined for C_{1-4} alkyl and pentyl, hexyl, 2-methylbutyl and the like; C_{2-6} alkyl as a group or part of a group defines straight or branched chain saturated hydrocarbon radicals having from 2 to 6 carbon atoms such as ethyl, propyl, 1-methylethyl, butyl, pentyl, hexyl, 2-methylbutyl and the like; C_{1-4} alkanediyl defines straight or branched chain saturated bivalent hydrocarbon radicals having from 1 to 4 carbon atoms such as methylene, 1,2-ethanediyl or 1,2-ethylidene, 1,3-propanediyl or 1,3-propylidene, 1,4-butanediyl or 1,4-butyldiene and the like; C_{3-7} cycloalkyl is generic to cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl and cycloheptyl; C_{2-6} alkenyl defines straight and branched chain hydrocarbon radicals having from 2 to 6 carbon atoms containing a double bond such as ethenyl, propenyl, butenyl, pentenyl, hexenyl and the like; C_{2-6} alkynyl defines straight and branched chain hydrocarbon radicals having from 2 to 6 carbon atoms containing a triple bond such as ethynyl, propynyl, butynyl, pentynyl, hexynyl and the like; a monocyclic, bicyclic or tricyclic saturated carbocycle represents a ring system consisting of 1, 2 or 3 rings, said ring system being composed of only carbon atoms and said ring system containing only single bonds; a monocyclic, bicyclic or tricyclic partially saturated carbocycle represents a ring system consisting of 1, 2 or 3 rings, said ring system being composed of only carbon atoms and comprising at least one double bond provided that the ring system is not an aromatic ring system; a monocyclic, bicyclic or tricyclic aromatic carbocycle represents an aromatic ring system consisting of 1, 2 or 3 rings, said ring system being composed of only carbon atoms; the term aromatic is well known to a person skilled in the art and designates cyclically conjugated systems of $4n + 2$ electrons, that is with 6, 10, 14 etc. π -electrons (rule of Hückel) ; a monocyclic, bicyclic or tricyclic saturated heterocycle represents a ring system consisting of 1, 2 or 3 rings and comprising at least one heteroatom selected

-10-

from O, N or S, said ring system containing only single bonds; a monocyclic, bicyclic or tricyclic partially saturated heterocycle represents a ring system consisting of 1, 2 or 3 rings and comprising at least one heteroatom selected from O, N or S, and at least one double bond provided that the ring system is not an aromatic ring system; a

5 monocyclic, bicyclic or tricyclic aromatic heterocycle represents an aromatic ring system consisting of 1, 2 or 3 rings and comprising at least one heteroatom selected from O, N or S.

Particular examples of monocyclic, bicyclic or tricyclic saturated carbocycles are
10 cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, cyclooctyl, bicyclo[4,2,0]octanyl, cyclononanyl, cyclodecanyl, decahydronaphthalenyl, tetradecahydroanthracenyl and the like.

Particular examples of monocyclic, bicyclic or tricyclic partially saturated carbocycles
15 are cyclopropenyl, cyclobutenyl, cyclopentenyl, cyclohexenyl, cycloheptenyl, cyclo-octenyl, bicyclo[4,2,0]octenyl, cyclononenyl, cyclodecenyl, octahydronaphthalenyl, 1,2,3,4-tetrahydronaphthalenyl, 1,2,3,4,4a,9,9a,10-octahydro-anthracenyl and the like.

Particular examples of monocyclic, bicyclic or tricyclic aromatic carbocycles are
20 phenyl, naphthalenyl, anthracenyl.

Particular examples of monocyclic, bicyclic or tricyclic saturated heterocycles are tetrahydrofuranyl, pyrrolidinyl, dioxolanyl, imidazolidinyl, thiazolidinyl, tetrahydrothienyl, dihydrooxazolyl, isothiazolidinyl, isoxazolidinyl, oxadiazolidinyl,
25 triazolidinyl, thiadiazolidinyl, pyrazolidinyl, piperidinyl, hexahydropyrimidinyl, hexahydropyrazinyl, dioxanyl, morpholinyl, dithianyl, thiomorpholinyl, piperazinyl, trithianyl, decahydroquinolinyl, octahydroindolyl and the like.

Particular examples of monocyclic, bicyclic or tricyclic partially saturated heterocycles
30 are pyrrolinyl, imidazolinyl, pyrazolinyl, 2,3-dihydrobenzofuranyl, 1,3-benzodioxolyl, 2,3-dihydro-1,4-benzodioxinyl, indolinyl and the like.

Particular examples of monocyclic, bicyclic or tricyclic aromatic heterocycles are azetyl, oxetylidenyl, pyrrolyl, furyl, thienyl, imidazolyl, oxazolyl, isoxazolyl, thiazolyl,
35 isothiazolyl, pyrazolyl, triazolyl, thiadiazolyl, oxadiazolyl, tetrazolyl, pyridyl, pyrimidinyl, pyrazinyl, pyridazinyl, triazinyl, pyranyl, benzofuryl, isobenzofuryl, benzothienyl, isobenzothienyl, indoliziny, indolyl, isoindolyl, benzoxazolyl,

-11-

benzimidazolyl, indazolyl, benzisoxazolyl, benzisothiazolyl, benzopyrazolyl,
benzoxadiazolyl, benzothiadiazolyl, benzotriazolyl, purinyl, quinolinyl, isoquinolinyl,
cinnolinyl, quinoliziny, phthalazinyl, quinoxaliny, quinazolinyl, naphthiridinyl,
pteridinyl, benzopyranyl, pyrrolopyridyl, thienopyridyl, furopyridyl, isothiazolopyridyl,
5 thiazolopyridyl, isoxazolopyridyl, oxazolopyridyl, pyrazolopyridyl, imidazopyridyl,
pyrrolopyrazinyl, thienopyrazinyl, furopyrazinyl, isothiazolopyrazinyl,
thiazolopyrazinyl, isoxazolopyrazinyl, oxazolopyrazinyl, pyrazolopyrazinyl,
imidazopyrazinyl, pyrrolopyrimidinyl, thienopyrimidinyl, fuopyrimidinyl,
isothiazolopyrimidinyl, thiazolopyrimidinyl, isoxazolopyrimidinyl,
10 oxazolopyrimidinyl, pyrazolopyrimidinyl, imidazopyrimidinyl, pyrrolopyridazinyl,
thienopyridazinyl, fuopyridazinyl, isothiazolopyridazinyl, thiazolopyridazinyl,
isoxazolopyridazinyl, oxazolopyridazinyl, pyrazolopyridazinyl, imidazopyridazinyl,
oxadiazolopyridyl, thiadiazolopyridyl, triazolopyridyl, oxadiazolopyrazinyl,
thiadiazolopyrazinyl, triazolopyrazinyl, oxadiazolopyrimidinyl, thiadiazolopyrimidinyl,
15 triazolopyrimidinyl, oxadiazolopyridazinyl, thiadiazolopyridazinyl, triazolopyridazinyl,
imidazooxazolyl, imidazothiazolyl, imidazoimidazolyl, isoxazolotriazinyl, isothiazolo-
triazinyl, pyrazolotriazinyl, oxazolotriazinyl, thiazolotriazinyl, imidazotriazinyl,
oxadiazolotriazinyl, thiadiazolotriazinyl, triazolotriazinyl, carbazolyl, acridinyl,
phenazinyl, phenothiazinyl, phenoxazinyl and the like.

20

As used herein before, the term (=O) forms a carbonyl moiety when attached to a carbon atom, a sulfoxide moiety when attached to a sulfur atom and a sulfonyl moiety when two of said terms are attached to a sulfur atom.

25

The term halo is generic to fluoro, chloro, bromo and iodo. As used in the foregoing and hereinafter, polyhaloC₁₋₄alkyl or polyhaloC₁₋₆alkyl as a group or part of a group is defined as mono- or polyhalosubstituted C₁₋₄alkyl or C₁₋₆alkyl, for example, as mono- or polyhalosubstituted methyl, in particular methyl with one or more fluoro atoms, for example, difluoromethyl or trifluoromethyl; 1,1-difluoro-ethyl and the like. In case
30 more than one halogen atoms are attached to an alkyl group within the definition of polyhaloC₁₋₄alkyl or polyhaloC₁₋₆alkyl, they may be the same or different.

35

The term heterocycle in the definition of R⁷ or R^{7a} is meant to include all the possible isomeric forms of the heterocycles, for instance, pyrrolyl comprises 1*H*-pyrrolyl and 2*H*-pyrrolyl.

-12-

The carbocycle or heterocycle in the definition of R^7 or R^{7a} may be attached to the remainder of the molecule of formula (I) through any ring carbon or heteroatom as appropriate, if not otherwise specified. Thus, for example, when the heterocycle is imidazolyl, it may be 1-imidazolyl, 2-imidazolyl, 4-imidazolyl and the like, or when
5 the carbocycle is naphthalenyl, it may be 1-naphthalenyl, 2-naphthalenyl and the like.

When any variable (eg. R^7 , X_2) occurs more than one time in any constituent, each definition is independent.

10 Lines drawn from substituents into ring systems indicate that the bond may be attached to any of the suitable ring atoms.

For therapeutic use, salts of the compounds of formula (I) are those wherein the counterion is pharmaceutically acceptable. However, salts of acids and bases which are
15 non-pharmaceutically acceptable may also find use, for example, in the preparation or purification of a pharmaceutically acceptable compound. All salts, whether pharmaceutically acceptable or not are included within the ambit of the present invention.

20 The pharmaceutically acceptable addition salts as mentioned hereinabove are meant to comprise the therapeutically active non-toxic acid addition salt forms which the compounds of formula (I) are able to form. The latter can conveniently be obtained by treating the base form with such appropriate acids as inorganic acids, for example, hydrohalic acids, e.g. hydrochloric, hydrobromic and the like; sulfuric acid; nitric acid;
25 phosphoric acid and the like; or organic acids, for example, acetic, propanoic, hydroxyacetic, 2-hydroxypropanoic, 2-oxopropanoic, oxalic, malonic, succinic, maleic, fumaric, malic, tartaric, 2-hydroxy-1,2,3-propanetricarboxylic, methanesulfonic, ethanesulfonic, benzenesulfonic, 4-methylbenzenesulfonic, cyclohexanesulfamic, 2-hydroxybenzoic, 4-amino-2-hydroxybenzoic and the like acids. Conversely the salt
30 form can be converted by treatment with alkali into the free base form.

The compounds of formula (I) containing acidic protons may be converted into their therapeutically active non-toxic metal or amine addition salt forms by treatment with appropriate organic and inorganic bases. Appropriate base salt forms comprise, for
35 example, the ammonium salts, the alkali and earth alkaline metal salts, e.g. the lithium, sodium, potassium, magnesium, calcium salts and the like, salts with organic bases, e.g. primary, secondary and tertiary aliphatic and aromatic amines such as methylamine,

-13-

ethylamine, propylamine, isopropylamine, the four butylamine isomers, dimethylamine, diethylamine, diethanolamine, dipropylamine, diisopropylamine, di-n-butylamine, pyrrolidine, piperidine, morpholine, trimethylamine, triethylamine, tripropylamine, quinuclidine, pyridine, quinoline and isoquinoline, the benzathine, 5 *N*-methyl-D-glucamine, 2-amino-2-(hydroxymethyl)-1,3-propanediol, hydrabamine salts, and salts with amino acids such as, for example, arginine, lysine and the like. Conversely the salt form can be converted by treatment with acid into the free acid form.

10 The term addition salt also comprises the hydrates and solvent addition forms which the compounds of formula (I) are able to form. Examples of such forms are e.g. hydrates, alcoholates and the like.

The term "quaternary amine" as used hereinbefore defines the quaternary ammonium salts which the compounds of formula (I) are able to form by reaction between a basic 15 nitrogen of a compound of formula (I) and an appropriate quaternizing agent, such as, for example, an optionally substituted alkylhalide, arylhalide or arylalkylhalide, e.g. methyl iodide or benzyl iodide. Other reactants with good leaving groups may also be used, such as alkyl trifluoromethanesulfonates, alkyl methanesulfonates, and alkyl *p*-toluenesulfonates. A quaternary amine has a positively charged nitrogen. 20 Pharmaceutically acceptable counterions include chloro, bromo, iodo, trifluoroacetate and acetate. The counterion of choice can be introduced using ion exchange resins.

The *N*-oxide forms of the present compounds are meant to comprise the compounds of formula (I) wherein one or several tertiary nitrogen atoms are oxidized to the so-called 25 *N*-oxide.

It will be appreciated that some of the compounds of formula (I) and their *N*-oxides, addition salts, quaternary amines and stereochemically isomeric forms may contain one or more centers of chirality and exist as stereochemically isomeric forms.

30 The term "stereochemically isomeric forms" as used hereinbefore defines all the possible stereoisomeric forms which the compounds of formula (I), and their *N*-oxides, addition salts, quaternary amines or physiologically functional derivatives may possess. Unless otherwise mentioned or indicated, the chemical designation of compounds 35 denotes the mixture of all possible stereochemically isomeric forms, said mixtures containing all diastereomers and enantiomers of the basic molecular structure as well as each of the individual isomeric forms of formula (I) and their *N*-oxides, salts, solvates


-14-

or quaternary amines substantially free, *i.e.* associated with less than 10%, preferably less than 5%, in particular less than 2% and most preferably less than 1% of the other isomers. Thus, when a compound of formula (I) is for instance specified as (E), this means that the compound is substantially free of the (Z) isomer.

- 5 In particular, stereogenic centers may have the R- or S-configuration; substituents on bivalent cyclic (partially) saturated radicals may have either the *cis*- or *trans*-configuration. Compounds encompassing double bonds can have an E (entgegen) or Z (zusammen) -stereochemistry at said double bond. The terms *cis*, *trans*, R, S, E and Z are well known to a person skilled in the art.
- 10 Stereochemically isomeric forms of the compounds of formula (I) are obviously intended to be embraced within the scope of this invention.


- Whenever used hereinbefore or hereinafter that substituents can be selected each independently out of a list of numerous definitions, such as for example for R⁹ and R¹⁰,
- 15 all possible combinations are intended which are chemically possible and which lead to chemically stable molecules.

- Some of the compounds of formula (I) may also exist in their tautomeric form. Such forms although not explicitly indicated in the above formula are intended to be included
- 20 within the scope of the present invention. For instance, the present invention includes the following compounds



- Whenever used hereinafter, the term "compounds of formula (I)" or any subgroup thereof, such as compounds of formula (I-A), (I-A-1), (I-A-2), (I-B), (I-B-1), (I-B-2), (I-C) or (I-D), is meant to also include their *N*-oxide forms, their salts, their quaternary amines and their stereochemically isomeric forms. Of special interest are those compounds of formula (I) which are stereochemically pure.
- 25
- 30 A first interesting embodiment of the present invention are those compounds of formula (I) having the following formula


-15-



a *N*-oxide, a pharmaceutically acceptable addition salt, a quaternary amine or a stereochemically isomeric form thereof, wherein R^1 , R^2 , R^3 , R^4 , ring E, ring F, C, D and m are as defined hereinabove.


5

A second interesting embodiment of the present invention are those compounds of formula (I-A) having the following formula



10 a *N*-oxide, a pharmaceutically acceptable addition salt, a quaternary amine or a stereochemically isomeric form thereof, wherein R^1 , R^2 , R^3 , R^4 , ring E, ring F, C, D and m are as defined hereinabove.

A third interesting embodiment of the present invention are those compounds of formula (I-A) having the following formula




15

a *N*-oxide, a pharmaceutically acceptable addition salt, a quaternary amine or a stereochemically isomeric form thereof, wherein R^1 , R^2 , R^3 , R^4 , ring E, ring F, C and D are as defined hereinabove.


-16-

A fourth interesting embodiment of the present invention are those compounds of formula (I) having the following formula



- 5 a *N*-oxide, a pharmaceutically acceptable addition salt, a quaternary amine or a stereochemically isomeric form thereof, wherein R^1 , R^2 , R^3 , R^4 , ring E, ring F, C, D and m are as defined hereinabove.


A fifth interesting embodiment of the present invention are those compounds of formula (I-B) having the following formula



10

- a *N*-oxide, a pharmaceutically acceptable addition salt, a quaternary amine or a stereochemically isomeric form thereof, wherein R^1 , R^2 , R^3 , R^4 , ring E, ring F, C, D and m are as defined hereinabove.

- 15 A sixth interesting embodiment of the present invention are those compounds of formula (I-B) having the following formula




20

- a *N*-oxide, a pharmaceutically acceptable addition salt, a quaternary amine or a stereochemically isomeric form thereof, wherein R^1 , R^2 , R^3 , R^4 , ring E, ring F, C and D are as defined hereinabove.


-17-

A seventh interesting embodiment of the present invention are those compounds of formula (I) having the following formula



- 5 a *N*-oxide, a pharmaceutically acceptable addition salt, a quaternary amine or a stereochemically isomeric form thereof, wherein A and B are as defined hereinabove and wherein R^{17a} represents C₁₋₆alkyl optionally substituted with hydroxy, cyano, aminocarbonyl, mono-or di(C₁₋₄alkyl)aminocarbonyl, C₁₋₄alkyloxycarbonyl or aryl.

- 10 An eighth interesting embodiment of the present invention are those compounds of formula (I) having the following formula



- 15 a *N*-oxide, a pharmaceutically acceptable addition salt, a quaternary amine or a stereochemically isomeric form thereof, wherein A and B are as defined hereinabove and wherein R^{17a} represents C₁₋₆alkyl optionally substituted with hydroxy, cyano, aminocarbonyl, mono-or di(C₁₋₄alkyl)aminocarbonyl, C₁₋₄alkyloxycarbonyl or aryl.

- A ninth interesting embodiment of the present invention are those compounds of formula (I), (I-A), (I-A-1), (I-A-2), (I-B), (I-B-1), (I-B-2), (I-C) or (I-D) wherein ring E represents phenyl.
- 20

- A tenth interesting embodiment of the present invention are those compounds of formula (I), (I-A), (I-A-1), (I-A-2), (I-B), (I-B-1), (I-B-2), (I-C) or (I-D) wherein ring F represents phenyl.
- 25

- A eleventh interesting embodiment of the present invention are those compounds of formula (I), (I-A), (I-A-1), (I-A-2), (I-B), (I-B-1), (I-B-2), (I-C) or (I-D) wherein ring E and ring F represent phenyl.

- 30 A twelfth interesting embodiment of the present invention are those compounds of formula (I) or any subgroup thereof as defined hereinabove wherein

-18-

- R^3 represents cyano; aminocarbonyl; amino; halo; NHR^{13} ; $NR^{13}R^{14}$; $-C(=O)-NHR^{13}$; $-C(=O)-NR^{13}R^{14}$; $-C(=O)-R^{15}$; $-CH=N-NH-C(=O)-R^{16}$; C_{1-6} alkyl optionally substituted with one or more substituents each independently selected from R^{3a} ; C_{1-6} alkyloxy optionally substituted with one or more substituents each independently selected from R^{3a} ; C_{1-6} alkyloxy C_{1-6} alkyl optionally substituted with one or more substituents each independently selected from R^{3a} ; C_{2-6} alkenyl optionally substituted with one or more substituents each independently selected from R^{3a} ; C_{2-6} alkynyl optionally substituted with one or more substituents each independently selected from R^{3a} ; or $-C(=N-O-R^8)-C_{1-4}$ alkyl;
- 10 R^{3a} represents halo, cyano, hydroxy, NR^9R^{10} , $-C(=O)-NR^9R^{10}$, $-C(=O)-C_{1-6}$ alkyl, $-C(=O)-O-C_{1-6}$ alkyl, $-C(=O)-polyhaloC_{1-6}$ alkyl, or $-C(=O)-O-polyhaloC_{1-6}$ alkyl; each R^4 independently represents hydroxy; halo; C_{1-6} alkyl optionally substituted with one or more substituents each independently selected from R^{4a} ; C_{2-6} alkenyl optionally substituted with one or more substituents each independently selected from R^{4a} ;
- 15 C_{2-6} alkynyl optionally substituted with one or more substituents each independently selected from R^{4a} ; C_{3-7} cycloalkyl; C_{1-6} alkyloxy; C_{1-6} alkyloxycarbonyl; C_{1-6} alkylcarbonyloxy; carboxyl; formyl; cyano; nitro; amino; mono- or di(C_{1-6} alkyl)amino; polyhalo C_{1-6} alkyl; polyhalo C_{1-6} alkyloxy; polyhalo C_{1-6} alkylthio; $-S(=O)_pR^6$; $-NH-S(=O)_pR^6$; $-C(=O)R^6$; $-NHC(=O)H$; $-C(=O)NHNH_2$; $NHC(=O)R^6$; or
- 20 $C(=NH)R^6$;
- R^9 and R^{10} each independently represent hydrogen; hydroxy; C_{1-6} alkyl; C_{1-6} alkyloxy; C_{1-6} alkylcarbonyl; C_{1-6} alkyloxycarbonyl; amino; mono- or di(C_{1-6} alkyl)amino; mono- or di(C_{1-6} alkyl)aminocarbonyl; $-CH(=NR^{11})$ or R^7 , wherein each of the aforementioned C_{1-6} alkyl groups may optionally and each individually be substituted
- 25 with one or two substituents each independently selected from hydroxy, C_{1-6} alkyloxy, hydroxy C_{1-6} alkyloxy, carboxyl, C_{1-6} alkyloxycarbonyl, cyano, amino, imino, mono- or di(C_{1-4} alkyl)amino, polyhalo C_{1-4} alkyl, polyhalo C_{1-4} alkyloxy, polyhalo C_{1-4} alkylthio, $-S(=O)_pR^6$, $-NH-S(=O)_pR^6$, $-C(=O)R^6$, $-NHC(=O)H$, $-C(=O)NHNH_2$, or $-NHC(=O)R^6$, $-C(=NH)R^6$, or
- 30 R^9 and R^{10} may be taken together to form a bivalent or trivalent radical of formula
- | | |
|------------------------------------|-----------|
| $-CH_2-CH_2-CH_2-CH_2-$ | (d-1); |
| $-CH_2-CH_2-CH_2-CH_2-CH_2-$ | (d-2); |
| $-CH_2-CH_2-O-CH_2-CH_2-$ | (d-3); |
| $-CH_2-CH_2-S-CH_2-CH_2-$ | (d-4); |
| 35 $-CH_2-CH_2-NR^{12}-CH_2-CH_2-$ | (d-5); |
| $-CH_2-CH=CH-CH_2-$ | (d-6); or |
| $=CH-CH=CH-CH=CH-$ | (d-7); |

-19-

R¹⁶ represents C₁₋₆alkyl optionally substituted with cyano, aminocarbonyl or mono- or di(C₁₋₄alkyl)aminocarbonyl;

aryl represents phenyl or phenyl substituted with one, two, three, four or five substituents each independently selected from halo, hydroxy, mercapto, C₁₋₆alkyl, hydroxyC₁₋₆alkyl, aminoC₁₋₆alkyl, mono or di(C₁₋₆alkyl)aminoC₁₋₆alkyl, C₁₋₆alkylcarbonyl, C₃₋₇cycloalkyl, C₁₋₆alkyloxy, C₁₋₆alkyloxycarbonyl, C₁₋₆alkylthio, cyano, nitro, polyhaloC₁₋₆alkyl, polyhaloC₁₋₆alkyloxy, or aminocarbonyl.

10 A thirteenth interesting embodiment of the present invention are those compounds of formula (I) or any subgroup thereof as defined hereinabove wherein ring E represents phenyl;

ring F represents phenyl;

R¹ represents hydrogen or C₁₋₆alkyl;

15 R² represents cyano; aminocarbonyl; mono- or di(C₁₋₄alkyl)aminocarbonyl; C₁₋₆alkyl substituted with cyano, aminocarbonyl or mono- or di(C₁₋₄alkyl)aminocarbonyl; or C₂₋₆alkenyl substituted with cyano, aminocarbonyl or mono- or di(C₁₋₄alkyl)aminocarbonyl; in particular cyano; C₁₋₆alkyl substituted with cyano, aminocarbonyl or mono- or di(C₁₋₄alkyl)aminocarbonyl; or C₂₋₆alkenyl substituted
20 with cyano, aminocarbonyl or mono- or di(C₁₋₄alkyl)aminocarbonyl; more in particular R² represents cyano or aminocarbonyl; more in particular R² represents cyano;

X₁ represents -NR⁵-; -O-; -C(=O)-; -C₁₋₄alkanediyl-; -CHOH-; -S-; or -S(=O)_p-; in particular X₁ represents -NR⁵-; -O- or -S-;

25 m represents an integer of value 1, 2 or 3; in particular 2;

R³ represents cyano; aminocarbonyl; amino; halo; NHR¹³; NR¹³R¹⁴; -C(=O)-NHR¹³; -C(=O)-NR¹³R¹⁴; -C(=O)-R¹⁵; C₁₋₆alkyl optionally substituted with one or more substituents each independently selected from R^{3a}; C₁₋₆alkyloxy optionally substituted with one or more substituents each independently selected from R^{3a};

30 C₁₋₆alkyloxyC₁₋₆alkyl optionally substituted with one or more substituents each independently selected from R^{3a}; C₂₋₆alkenyl optionally substituted with one or more substituents each independently selected from R^{3a}; or C₂₋₆alkynyl optionally substituted with one or more substituents each independently selected from R^{3a}; in particular R³ represents cyano; aminocarbonyl; C₁₋₆alkyl optionally substituted with
35 one or more substituents each independently selected from R^{3a}; C₁₋₆alkyloxy optionally substituted with one or more substituents each independently selected from

-20-

- R^{3a} ; or C₂₋₆alkenyl optionally substituted with one or more substituents each independently selected from R^{3a} ;
- R^{3a} represents halo, cyano, hydroxy, NR^9R^{10} , $-C(=O)-NR^9R^{10}$, $-C(=O)-C_{1-6}$ alkyl, $-C(=O)-O-C_{1-6}$ alkyl, $-C(=O)-polyhaloC_{1-6}$ alkyl, or $-C(=O)-O-polyhaloC_{1-6}$ alkyl; in particular R^{3a} represents cyano or aminocarbonyl;
- each R^4 independently represents hydroxy; halo; C₁₋₆alkyl optionally substituted with one or more substituents each independently selected from R^{4a} ; C₂₋₆alkenyl optionally substituted with one or more substituents each independently selected from R^{4a} ; C₂₋₆alkynyl optionally substituted with one or more substituents each independently selected from R^{4a} ; C₃₋₇cycloalkyl; C₁₋₆alkyloxy; C₁₋₆alkyloxycarbonyl; C₁₋₆alkylcarbonyloxy; carboxyl; formyl; cyano; nitro; amino; mono- or di(C₁₋₆alkyl)amino; polyhaloC₁₋₆alkyl; polyhaloC₁₋₆alkyloxy; polyhaloC₁₋₆alkylthio; $-S(=O)_pR^6$; $-NH-S(=O)_pR^6$; or $-C(=O)R^6$; in particular R^4 represents halo; C₁₋₆alkyl; C₁₋₆alkyloxy; cyano; amino; polyhaloC₁₋₆alkyl; or polyhaloC₁₋₆alkyloxy;
- R^{4a} represents halo, cyano, NR^9R^{10} , hydroxy or $-C(=O)R^6$;
- R^5 represents hydrogen or C₁₋₆alkyl;
- R^9 and R^{10} each independently represent hydrogen; hydroxy; C₁₋₆alkyl; C₁₋₆alkyloxy; C₁₋₆alkylcarbonyl; C₁₋₆alkyloxycarbonyl; amino; mono- or di(C₁₋₆alkyl)amino; or mono- or di(C₁₋₆alkyl)aminocarbonyl; wherein each of the aforementioned C₁₋₆alkyl groups may optionally and each individually be substituted with one or two substituents each independently selected from hydroxy, C₁₋₆alkyloxy, hydroxyC₁₋₆alkyloxy, carboxyl, C₁₋₆alkyloxycarbonyl, cyano, amino, imino, mono- or di(C₁₋₄alkyl)amino, polyhaloC₁₋₄alkyl, polyhaloC₁₋₄alkyloxy, polyhaloC₁₋₄alkylthio, $-S(=O)_pR^6$, $-NH-S(=O)_pR^6$, or $-C(=O)R^6$; or
- R^9 and R^{10} may be taken together to form a bivalent or trivalent radical of formula
- | | |
|---------------------------------|----------------------------------------------------------------------------------------------------------------------------------------------|
| $-CH_2-CH_2-CH_2-CH_2-$ | (d-1); |
| $-CH_2-CH_2-CH_2-CH_2-CH_2-$ | (d-2); |
| $-CH_2-CH_2-O-CH_2-CH_2-$ | (d-3); |
| $-CH_2-CH_2-S-CH_2-CH_2-$ | (d-4); |
| $-CH_2-CH_2-NR^{12}-CH_2-CH_2-$ | (d-5); |
| $-CH_2-CH=CH-CH_2-$ | (d-6); or |
| $=CH-CH=CH-CH=CH-$ | (d-7); in particular R^9 and R^{10} each independently represent hydrogen; hydroxy; C ₁₋₆ alkyl or C ₁₋₆ alkyloxy; |
- aryl represents phenyl or phenyl substituted with one, two, three, four or five substituents each independently selected from halo, hydroxy, mercapto, C₁₋₆alkyl, hydroxyC₁₋₆alkyl, aminoC₁₋₆alkyl, mono or di(C₁₋₆alkyl)aminoC₁₋₆alkyl, C₁₋₆alkylcarbonyl, C₃₋₇cycloalkyl, C₁₋₆alkyloxy, C₁₋₆alkyloxycarbonyl,

-21-

C₁₋₆alkylthio, cyano, nitro, polyhaloC₁₋₆alkyl, polyhaloC₁₋₆alkyloxy, or aminocarbonyl.

5 A fourteenth interesting embodiment of the present invention are those compounds of formula (I) or any subgroup thereof as defined hereinabove, wherein, whenever the substituents are present, m is 1, 2 or 3; in particular 2.

10 A fifteenth interesting embodiment of the present invention are those compounds of formula (I) or any subgroup thereof as defined hereinabove, wherein R¹ is hydrogen or C₁₋₆alkyl; in particular hydrogen.

15 A sixteenth interesting embodiment of the present invention are those compounds of formula (I) or any subgroup thereof as defined hereinabove wherein X₁ represents -NR⁵-; -O-; -C(=O)-; -C₁₋₄alkanediyl-; -CHOH-; -S-; or -S(=O)_p-; in particular X₁ represents -NR⁵-; -O-; or -S-; with R⁵ representing hydrogen or C₁₋₆alkyl.


20 A seventeenth interesting embodiment of the present invention are those compounds of formula or any subgroup thereof as defined hereinabove wherein m is 2 and R³ represents cyano; C₁₋₆alkyl optionally substituted with cyano; C₁₋₆alkyloxy optionally substituted with cyano; C₂₋₆alkenyl substituted with cyano or -C(=O)-NR⁹R¹⁰.

25 An eighteenth interesting embodiment of the present invention are those compounds of formula (I) as defined above wherein R² represents cyano; aminocarbonyl; mono- or di(C₁₋₄alkyl)aminocarbonyl; C₁₋₆alkyl substituted with cyano, aminocarbonyl or mono- or di(C₁₋₄alkyl)aminocarbonyl; C₂₋₆alkenyl substituted with cyano, aminocarbonyl or mono- or di(C₁₋₄alkyl)aminocarbonyl; or C₂₋₆alkynyl substituted with cyano, aminocarbonyl or mono- or di(C₁₋₄alkyl)aminocarbonyl.

30 A nineteenth interesting embodiment of the present invention are those compounds of formula (I) as defined above wherein R² represents cyano or aminocarbonyl; in particular cyano.

35 An twentieth interesting embodiment of the present invention is the use of a compound of formula (I') for the manufacture of a medicament for the prevention or the treatment of HIV infection, wherein the compound of formula (I') is a compound of formula

-22-



a *N*-oxide, a pharmaceutically acceptable addition salt, a quaternary amine or a stereochemically isomeric form thereof, wherein

$-a^1=a^2-C(R^2)=a^3-a^4-$ represents a bivalent radical of formula

5 $-CH=CH-C(R^2)=CH-CH=$ (a-1);

$-N=CH-C(R^2)=CH-CH=$ (a-2);

$-CH=N-C(R^2)=CH-CH=$ (a-3);

$-N=CH-C(R^2)=N-CH=$ (a-4);

$-N=CH-C(R^2)=CH-N=$ (a-5);

10 $-CH=N-C(R^2)=N-CH=$ (a-6); or

$-N=N-C(R^2)=CH-CH=$ (a-7);

$-b^1=b^2-b^3=b^4-$ represents a bivalent radical of formula

$-CH=CH-CH=CH-$ (b-1);

$-N=CH-CH=CH-$ (b-2);

15 $-N=CH-N=CH-$ (b-3);

$-N=CH-CH=N-$ (b-4); or

$-N=N-CH=CH-$ (b-5);

$-C-D-$ represents a bivalent radical of formula

$-N=CH-NR^{17}-$ (c-1); or

20 $-NR^{17}-CH=N-$ (c-2);

m represents an integer of value 1, 2, 3 and in case $-b^1=b^2-b^3=b^4-$ is (b-1), then m may also be 4;

R^1 represents hydrogen; aryl; formyl; C_{1-6} alkylcarbonyl; C_{1-6} alkyloxycarbonyl;

C_{1-6} alkyl optionally substituted with formyl, C_{1-6} alkylcarbonyl,

25 C_{1-6} alkyloxycarbonyl, C_{1-6} alkylcarbonyloxy; or C_{1-6} alkyloxy C_{1-6} alkylcarbonyl substituted with C_{1-6} alkyloxycarbonyl;

R^2 represents cyano; aminocarbonyl; mono- or di(C_{1-4} alkyl)aminocarbonyl; C_{1-6} alkyl optionally substituted with cyano, aminocarbonyl or mono- or di(C_{1-4} alkyl)aminocarbonyl; C_{2-6} alkenyl substituted with cyano, aminocarbonyl or

-23-

- mono- or di(C₁₋₄alkyl)aminocarbonyl; or C₂₋₆alkynyl substituted with cyano, aminocarbonyl or mono- or di(C₁₋₄alkyl)aminocarbonyl;
- X₁ represents -NR⁵-, -NH-NH-, -N=N-, -O-, -C(=O)-, C₁₋₄alkanediyl, -CHOH-, -S-, -S(=O)_p-, -X₂-C₁₋₄alkanediyl- or -C₁₋₄alkanediyl-X₂-;
- 5 X₂ represents -NR⁵-, -NH-NH-, -N=N-, -O-, -C(=O)-, -CHOH-, -S-, -S(=O)_p-;
- R³ represents NHR¹³; NR¹³R¹⁴; -C(=O)-NHR¹³; -C(=O)-NR¹³R¹⁴; -C(=O)-R¹⁵; -CH=N-NH-C(=O)-R¹⁶; cyano; halo; C₁₋₆alkyl; polyhaloC₁₋₆alkyl; C₁₋₆alkyl substituted with one or more substituents each independently selected from cyano, NR⁹R¹⁰, -C(=O)-NR⁹R¹⁰, -C(=O)-C₁₋₆alkyl or R⁷; C₁₋₆alkyl substituted with hydroxy and a second substituent selected from cyano, NR⁹R¹⁰, -C(=O)-NR⁹R¹⁰, -C(=O)-C₁₋₆alkyl or R⁷; C₁₋₆alkyloxyC₁₋₆alkyl optionally substituted with one or more substituents each independently selected from cyano, NR⁹R¹⁰, -C(=O)-NR⁹R¹⁰, -C(=O)-C₁₋₆alkyl or R⁷; C₁₋₆alkyloxy optionally substituted with one or more substituents each independently selected from cyano, NR⁹R¹⁰, -C(=O)-NR⁹R¹⁰, -C(=O)-C₁₋₆alkyl or R⁷; C₂₋₆alkenyl optionally substituted with one or more substituents each independently selected from halo, cyano, NR⁹R¹⁰, -C(=O)-NR⁹R¹⁰, -C(=O)-C₁₋₆alkyl or R⁷; C₂₋₆alkynyl optionally substituted with one or more substituents each independently selected from halo, cyano, NR⁹R¹⁰, -C(=O)-NR⁹R¹⁰, -C(=O)-C₁₋₆alkyl or R⁷; -C(=N-O-R⁸)-C₁₋₄alkyl; R⁷ or -X₃-R⁷;
- 10 X₃ is -NR⁵-, -NH-NH-, -N=N-, -O-, -C(=O)-, -S-, -S(=O)_p-, -X_{4b}-C₁₋₄alkanediyl-, -C₁₋₄alkanediyl-X_{4a}-, -C₁₋₄alkanediyl-X_{4b}-C₁₋₄alkanediyl-, -C(=N-OR⁸)-C₁₋₄alkanediyl-;
- 15 with X_{4a} being -NH-NH-, -N=N-, -O-, -C(=O)-, -S-, -S(=O)_p-; and with X_{4b} being -NH-NH-, -N=N-, -C(=O)-, -S-, -S(=O)_p-;
- 20 each R⁴ independently represents halo, hydroxy, C₁₋₆alkyl, C₃₋₇cycloalkyl, C₁₋₆alkyloxy, hydroxyC₁₋₆alkyl, aminoC₁₋₆alkyl, cyano, nitro, polyhaloC₁₋₆alkyl, polyhaloC₁₋₆alkyloxy, aminocarbonyl, mono- or di(C₁₋₄alkyl)aminocarbonyl, C₁₋₆alkyloxycarbonyl, C₁₋₆alkylcarbonyl, formyl, amino, mono- or di(C₁₋₄alkyl)amino or R⁷;
- 25 R⁵ is hydrogen; aryl; formyl; C₁₋₆alkylcarbonyl; C₁₋₆alkyloxycarbonyl; C₁₋₆alkyl optionally substituted with formyl, C₁₋₆alkylcarbonyl, C₁₋₆alkyloxycarbonyl or C₁₋₆alkylcarbonyloxy; or C₁₋₆alkyloxyC₁₋₆alkylcarbonyl substituted with C₁₋₆alkyloxycarbonyl;
- 30 R⁶ is C₁₋₄alkyl, amino, mono- or di(C₁₋₄alkyl)amino or polyhaloC₁₋₄alkyl;
- 35 R⁷ is a monocyclic, bicyclic or tricyclic saturated, partially saturated or aromatic carbocycle or a monocyclic, bicyclic or tricyclic saturated, partially saturated or aromatic heterocycle, wherein each of said carbocyclic or heterocyclic ring systems

-24-

- may optionally be substituted where possible with one, two, three, four or five substituents each independently selected from halo, hydroxy, mercapto, C₁₋₆alkyl, hydroxyC₁₋₆alkyl, aminoC₁₋₆alkyl, mono or di(C₁₋₆alkyl)aminoC₁₋₆alkyl, formyl, C₁₋₆alkylcarbonyl, C₃₋₇cycloalkyl, C₁₋₆alkyloxy, C₁₋₆alkyloxycarbonyl, C₁₋₆alkylthio, cyano, nitro, polyhaloC₁₋₆alkyl, polyhaloC₁₋₆alkyloxy, aminocarbonyl, -CH(=N-O-R⁸), R^{7a}, -X₃-R^{7a} or R^{7a}-C₁₋₄alkanediyl-;
- 5 R^{7a} is a monocyclic, bicyclic or tricyclic saturated, partially saturated or aromatic carbocycle or a monocyclic, bicyclic or tricyclic saturated, partially saturated or aromatic heterocycle, wherein each of said carbocyclic or heterocyclic ring systems may optionally be substituted where possible with one, two, three, four or five substituents each independently selected from halo, hydroxy, mercapto, C₁₋₆alkyl, hydroxyC₁₋₆alkyl, aminoC₁₋₆alkyl, mono or di(C₁₋₆alkyl)aminoC₁₋₆alkyl, formyl, C₁₋₆alkylcarbonyl, C₃₋₇cycloalkyl, C₁₋₆alkyloxy, C₁₋₆alkyloxycarbonyl, C₁₋₆alkylthio, cyano, nitro, polyhaloC₁₋₆alkyl, polyhaloC₁₋₆alkyloxy, aminocarbonyl, or -CH(=N-O-R⁸);
- 10 R⁸ is hydrogen, C₁₋₄alkyl optionally substituted with aryl, or aryl;
- 15 R⁹ and R¹⁰ each independently are hydrogen; C₁₋₆alkyl; C₁₋₆alkylcarbonyl; C₁₋₆alkyloxycarbonyl; amino; mono- or di(C₁₋₆alkyl)amino; mono- or di(C₁₋₆alkyl)aminocarbonyl; -CH(=NR¹¹) or R⁷, wherein each of the aforementioned C₁₋₆alkyl groups may optionally and each individually be substituted with one or two substituents each independently selected from hydroxy, C₁₋₆alkyloxy, hydroxyC₁₋₆alkyloxy, carboxyl, C₁₋₆alkyloxycarbonyl, cyano, amino, imino, mono- or di(C₁₋₄alkyl)amino, polyhaloC₁₋₄alkyl, polyhaloC₁₋₄alkyloxy, polyhaloC₁₋₄alkylthio, -S(=O)_pR⁶, -NH-S(=O)_pR⁶, -C(=O)R⁶, -NHC(=O)H, -C(=O)NHNH₂, -NHC(=O)R⁶, -C(=NH)R⁶, R⁷; or
- 20 R⁹ and R¹⁰ may be taken together to form a bivalent or trivalent radical of formula
- 25 -CH₂-CH₂-CH₂-CH₂- (d-1);
- CH₂-CH₂-CH₂-CH₂-CH₂- (d-2);
- CH₂-CH₂-O-CH₂-CH₂- (d-3);
- 30 -CH₂-CH₂-S-CH₂-CH₂- (d-4);
- CH₂-CH₂-NR¹²-CH₂-CH₂- (d-5);
- CH₂-CH=CH-CH₂- (d-6); or
- =CH-CH=CH-CH=CH- (d-7);
- R¹¹ represents cyano; C₁₋₄alkyl optionally substituted with C₁₋₄alkyloxy, cyano, amino, mono- or di(C₁₋₄alkyl)amino or aminocarbonyl; C₁₋₄alkylcarbonyl; C₁₋₄alkyloxycarbonyl; aminocarbonyl; mono- or di(C₁₋₄alkyl)aminocarbonyl;
- 35 R¹² represents hydrogen or C₁₋₄alkyl;

-25-

- R^{13} and R^{14} each independently represent C_{1-6} alkyl optionally substituted with cyano, aminocarbonyl or mono- or di(C_{1-4} alkyl)aminocarbonyl; C_{2-6} alkenyl optionally substituted with cyano, aminocarbonyl or mono- or di(C_{1-4} alkyl)aminocarbonyl; C_{2-6} alkynyl optionally substituted with cyano, aminocarbonyl or mono- or di(C_{1-4} alkyl)aminocarbonyl;
- 5 R^{15} represents C_{1-6} alkyl substituted with cyano, aminocarbonyl or mono- or di(C_{1-4} alkyl)aminocarbonyl;
- R^{16} represents C_{1-6} alkyl optionally substituted with cyano, aminocarbonyl or mono- or di(C_{1-4} alkyl)aminocarbonyl; or R^7 ;
- 10 R^{17} represents hydrogen; C_{1-6} alkyl; or C_{1-6} alkyl substituted with aryl;
- p is 1 or 2;
- aryl represents phenyl or phenyl substituted with one, two, three, four or five substituents each independently selected from halo, hydroxy, mercapto, C_{1-6} alkyl, hydroxy C_{1-6} alkyl, amino C_{1-6} alkyl, mono or di(C_{1-6} alkyl)amino C_{1-6} alkyl,
- 15 C_{1-6} alkylcarbonyl, C_{3-7} cycloalkyl, C_{1-6} alkyloxy, C_{1-6} alkyloxy carbonyl, C_{1-6} alkylthio, cyano, nitro, polyhalo C_{1-6} alkyl, polyhalo C_{1-6} alkyloxy, aminocarbonyl, R^7 or $-X_3-R^7$.

- A twenty first interesting embodiment of the present invention are those compounds of formula (I') as defined hereinabove provided that when R^2 represents aminocarbonyl or mono- or di(C_{1-4} alkyl)aminocarbonyl then R^3 represents $-C(=O)-R^{15}$; $-\text{CH}=\text{N}-\text{NH}-C(=O)-R^{16}$; cyano; C_{1-6} alkyl substituted with one or more substituents each independently selected from cyano, NR^9R^{10} , $-C(=O)-\text{NR}^9\text{R}^{10}$, $-C(=O)-C_{1-6}$ alkyl or R^7 ; C_{1-6} alkyl substituted with hydroxy and a second substituent selected from cyano, NR^9R^{10} , $-C(=O)-\text{NR}^9\text{R}^{10}$, $-C(=O)-C_{1-6}$ alkyl or R^7 ; C_{1-6} alkyloxy C_{1-6} alkyl optionally substituted with one or more substituents each independently selected from cyano, NR^9R^{10} , $-C(=O)-\text{NR}^9\text{R}^{10}$, $-C(=O)-C_{1-6}$ alkyl or R^7 ; C_{1-6} alkyloxy substituted with one or more substituents each independently selected from cyano, NR^9R^{10} , $-C(=O)-\text{NR}^9\text{R}^{10}$, $-C(=O)-C_{1-6}$ alkyl or R^7 ; C_{2-6} alkenyl optionally substituted with one or more substituents each independently selected from halo, cyano, NR^9R^{10} , $-C(=O)-\text{NR}^9\text{R}^{10}$, $-C(=O)-C_{1-6}$ alkyl or R^7 ; C_{2-6} alkynyl optionally substituted with one or more substituents each independently selected from halo, cyano, NR^9R^{10} , $-C(=O)-\text{NR}^9\text{R}^{10}$, $-C(=O)-C_{1-6}$ alkyl or R^7 ; $-C(=\text{N}-\text{O}-\text{R}^8)-C_{1-4}$ alkyl; R^7 or $-X_3-R^7$.
- 20
- 25
- 30
- 35 A twenty second interesting embodiment of the present invention are those compounds of formula (I) or any subgroup thereof as defined hereinabove wherein R^3 is R^7 , $\text{NR}^{13}\text{R}^{14}$, $-C(=O)\text{R}^{15}$, $-\text{CH}=\text{N}-\text{NH}-C(=O)\text{R}^{16}$, $-C(=O)\text{NHR}^{13}$, $-C(=O)\text{NR}^{13}\text{R}^{14}$,

-26-

- C(=N-OR⁸)-C₁₋₄alkyl, C₁₋₆alkyl substituted with cyano, C₁₋₆alkyl substituted twice with cyano, C₁₋₆alkyl substituted with NR⁹R¹⁰, C₁₋₆alkyl substituted with hydroxy and cyano, C₁₋₆alkyl substituted with hydroxy and R⁷, C₁₋₆alkyloxy C₁₋₆alkyl, C₁₋₆alkyloxy C₁₋₆alkyl substituted with cyano, C₂₋₆alkenyl substituted with R⁷, C₂₋₆alkenyl substituted with cyano, C₂₋₆alkenyl substituted twice with cyano, C₂₋₆alkenyl substituted with cyano and R⁷, C₂₋₆alkenyl substituted with cyano and -C(=O)-C₁₋₆alkyl, C₂₋₆alkenyl substituted with cyano and halo, C₂₋₆alkenyl substituted with -C(=O)-NR⁹R¹⁰, C₂₋₆alkenyl substituted with halo, C₂₋₆alkenyl substituted twice with halo or C₂₋₆alkenyl substituted with NR⁹R¹⁰.
- 5 substituted with cyano, C₂₋₆alkenyl substituted twice with cyano, C₂₋₆alkenyl substituted with cyano and R⁷, C₂₋₆alkenyl substituted with cyano and -C(=O)-C₁₋₆alkyl, C₂₋₆alkenyl substituted with cyano and halo, C₂₋₆alkenyl substituted with -C(=O)-NR⁹R¹⁰, C₂₋₆alkenyl substituted with halo, C₂₋₆alkenyl substituted twice with halo or C₂₋₆alkenyl substituted with NR⁹R¹⁰.
- 10 A twenty third interesting embodiment of the present invention are those compounds of formula (I) or any subgroup thereof as defined hereinabove wherein R³ is cyano; aminocarbonyl; C₁₋₆alkyl optionally substituted with cyano or aminocarbonyl; C₁₋₆alkyloxy optionally substituted with cyano or aminocarbonyl; C₂₋₆alkenyl substituted with cyano or aminocarbonyl.
- 15 A twenty fourth interesting embodiment of the present invention are those compounds of formula (I) or any subgroup thereof as defined hereinabove wherein R³ represents cyano; C₁₋₆alkyl; C₁₋₆alkyl substituted with cyano; C₁₋₆alkyloxy optionally substituted with cyano; C₂₋₆alkenyl substituted with cyano or -C(=O)-NR⁹R¹⁰;
- 20 A twenty fifth interesting embodiment of the present invention are those compounds of formula (I) or any subgroup thereof as defined hereinabove wherein m is 2 and each R⁴ independently represents halo, C₁₋₆alkyl or C₁₋₆alkyloxy.
- 25 A twenty sixth interesting embodiment of the present invention are those compounds of formula (I) or any subgroup thereof as defined hereinabove wherein R¹⁷ represents hydrogen.
- 30 A twenty seventh interesting embodiment of the present invention are those compounds of formula (I) as defined hereinabove wherein R¹⁷ represents C₁₋₆alkyl or C₁₋₆alkyl substituted with hydroxy, cyano or aryl.
- 35 A twenty eighth interesting embodiment of the present invention are those compounds of formula (I) or any subgroup thereof as defined hereinabove wherein ring F is 2,4,6-trisubstituted phenyl.

-27-

A twenty ninth interesting embodiment of the present invention are those compounds of formula (I) or any subgroup thereof as defined hereinabove wherein the following definitions, apply :

- ring E represents phenyl;
5 ring F represents phenyl;
m is 2;
R¹ represents hydrogen;
R² represents cyano, aminocarbonyl or C₁₋₆alkyl, in particular cyano or
C₁₋₆alkyl; more in particular cyano;
10 R³ represents cyano; C₁₋₆alkyl; C₁₋₆alkyl substituted with cyano; C₁₋₆alkyloxy
optionally substituted with cyano; C₂₋₆alkenyl substituted with cyano or
-C(=O)-NR⁹R¹⁰;
each R⁴ independently represents halo, C₁₋₆alkyl or C₁₋₆alkyloxy;
X₁ represents -NR⁵- or -O-;
15 R⁵ represents hydrogen;
R⁹ and R¹⁰ each independently are hydrogen or C₁₋₆alkyl; or R⁹ and R¹⁰ may be taken
together to form a bivalent radical of formula -CH₂-CH₂-O-CH₂-CH₂- (d-3);
R¹⁷ is hydrogen; C₁₋₆alkyl optionally substituted with hydroxy, cyano, aminocarbonyl,
C₁₋₄alkyloxycarbonyl or aryl;
20 aryl is phenyl substituted with C₁₋₆alkyloxy.


A thirtieth interesting embodiment of the present invention are those compounds of formula (I-A) or (I-B) as defined hereinabove wherein the following definitions, apply :

- R¹ is hydrogen;
25 R² is cyano or aminocarbonyl, in particular cyano;
R³ is C₂₋₆alkenyl substituted with cyano; C₁₋₆alkyloxy optionally substituted with
cyano; cyano; C₁₋₆alkyl optionally substituted with cyano;
R⁴ is C₁₋₆alkyl, halo, C₁₋₆alkyloxy;
X₁ is NH or O;
30 ring E is phenyl;
ring F is phenyl;
m is 2.

- Preferred compounds of formula (I) are compounds 13, 26, 19, 54, 55, 45, 46, 72, 64,
35 53, 62, 52, 48, 50, 63, 49, 34, 2, 6, 31, 32, 73, 70 and 47, (see Tables 1, 2, 3 and 4),
their N-oxides, pharmaceutically acceptable addition salts, quaternary amines and
stereochemically isomeric forms thereof.


-28-

- In general, compounds of formula (I) wherein R^{17} represents C_{1-6} alkyl optionally substituted with aryl, said R^{17} being represented by R^{17b} , and said compounds being represented by formula (I-a) or (I-b), can be prepared by reacting an intermediate of formula (II-a) or (II-b) wherein W_1 represents a suitable leaving group, such as for example halogen, e.g. chloro and the like, with an intermediate of formula (III) in the presence of a suitable catalyst, such as for example $Pd(OAc)_2$, $Pd_2(dba)_3$ and the like, a suitable ligand, such as for example a mixture of (+) and (-) 2,2'-bis(diphenylphosphino)-1,1'-binaphthyl, a suitable base, such as for example Cs_2CO_3 or Na_3PO_4 , and a suitable solvent, such as for example toluene or dioxane.



- Compounds of formula (I-a) or (I-b) may also be prepared by reacting an intermediate of formula (II'-a) or (II'-b) wherein W_1 represents a suitable leaving group, such as for example halogen, e.g. chloro and the like, with an intermediate of formula (III') in the presence of a suitable catalyst, such as for example $Pd(OAc)_2$, $Pd_2(dba)_3$ and the like, a suitable ligand, such as for example a mixture of (+) and (-) 2,2'-bis(diphenylphosphino)-1,1'-binaphthyl, a suitable base, such as for example Cs_2CO_3 or Na_3PO_4 , and a suitable solvent, such as for example toluene or dioxane.

-29-



The compounds of formula (I) may further be prepared by converting compounds of formula (I) into each other according to art-known group transformation reactions.

- 5 The compounds of formula (I) may be converted to the corresponding *N*-oxide forms following art-known procedures for converting a trivalent nitrogen into its *N*-oxide form. Said *N*-oxidation reaction may generally be carried out by reacting the starting material of formula (I) with an appropriate organic or inorganic peroxide. Appropriate inorganic peroxides comprise, for example, hydrogen peroxide, alkali metal or earth
- 10 alkaline metal peroxides, e.g. sodium peroxide, potassium peroxide; appropriate organic peroxides may comprise peroxy acids such as, for example, benzenecarboperoxoic acid or halo substituted benzenecarboperoxoic acid, e.g. 3-chlorobenzenecarboperoxoic acid, peroxyalkanoic acids, e.g. peroxyacetic acid, alkylhydroperoxides, e.g. tert.butyl hydro-peroxide. Suitable solvents are, for example, water, lower alcohols,
- 15 e.g. ethanol and the like, hydrocarbons, e.g. toluene, ketones, e.g. 2-butanone, halogenated hydrocarbons, e.g. dichloromethane, and mixtures of such solvents.

- For instance, a compound of formula (I) wherein R³ comprises cyano, can be converted into a compound of formula (I) wherein R³ comprises aminocarbonyl, by reaction with
- 20 HCOOH, in the presence of a suitable acid, such as hydrochloric acid. A compound of formula (I) wherein R³ comprises cyano, can also further be converted into a compound of formula (I) wherein R³ comprises tetrazolyl, by reaction with sodium azide in the presence of ammonium chloride and *N,N*-dimethylacetamide.

-30-

Compounds of formula (I) wherein R^3 comprises aminocarbonyl, can be converted into a compound of formula (I) wherein R^3 comprises cyano, in the presence of a suitable dehydrating agent. The dehydration can be performed according to methodologies well-known to the person skilled in the art, such as the ones disclosed in

- 5 "Comprehensive Organic Transformations. A guide to functional group preparations" by Richard C. Larock, John Wiley & Sons, Inc, 1999, p 1983-1985, which is incorporated herein by reference. Different suitable reagents are enumerated in said reference, such as for example SOCl_2 , HOSO_2NH_2 , ClSO_2NCO , $\text{MeO}_2\text{CNSO}_2\text{NEt}_3$, PhSO_2Cl , TsCl , P_2O_5 , $(\text{Ph}_3\text{PO}_3\text{SCF}_3)_3\text{O}_3\text{SCF}_3$, polyphosphate ester, $(\text{EtO})_2\text{POP}(\text{OEt})_2$,
10 $(\text{EtO})_3\text{PI}_2$, 2-chloro-1,3,2-dioxaphospholane, 2,2,2-trichloro-2,2-dihydro-1,3,2-dioxaphospholane, POCl_3 , PPh_3 , $\text{P}(\text{NCl}_2)_3$, $\text{P}(\text{NEt}_2)_3$, COCl_2 , $\text{NaCl} \cdot \text{AlCl}_3$, ClCOCOCl , ClCO_2Me , Cl_3CCOCl , $(\text{CF}_3\text{CO})_2\text{O}$, $\text{Cl}_3\text{CN}=\text{CCl}_2$, 2,4,6-trichloro-1,3,5-triazine, $\text{NaCl} \cdot \text{AlCl}_3$, $\text{HN}(\text{SiMe}_2)_3$, $\text{N}(\text{SiMe}_2)_4$, LiAlH_4 and the like. All the reagents listed in said publication are incorporated herein by reference.

15

Compounds of formula (I) wherein R^3 represents optionally substituted C_{2-6} alkenyl, can be converted into a compound of formula (I) wherein R^3 represents optionally substituted C_{1-6} alkyl, by reduction in the presence of a suitable reducing agent, such as for example H_2 , in the presence of a suitable catalyst, such as for example palladium on
20 charcoal, and in the presence of a suitable solvent, such as for example an alcohol, e.g. methanol.

20

- Compounds of formula (I) wherein R^3 represents $\text{CH}(\text{OH})\text{-R}^{16}$, can be converted into a compound of formula (I) wherein R^3 represents $\text{C}(=\text{O})\text{-R}^{16}$ by reaction with Jones's
25 reagent in the presence of a suitable solvent, such as for example 2-propanone.

25

- Compound of formula (I) wherein R^3 represents $\text{C}(=\text{O})\text{-CH}_2\text{-R}^{16a}$, wherein R^{16a} represents cyano, aminocarbonyl or mono -or di(C_{1-4} alkyl)aminocarbonyl, can be converted into a compound of formula (I) wherein R^3 represents $\text{C}(\text{Cl})=\text{CH-R}^{16a}$ by
30 reaction with POCl_3 .

30

- Compounds of formula (I) wherein R^3 represents a monocyclic, bicyclic or tricyclic saturated carbocycle; a monocyclic, bicyclic or tricyclic partially saturated carbocycle; a monocyclic, bicyclic or tricyclic aromatic carbocycle; a monocyclic, bicyclic or
35 tricyclic saturated heterocycle; a monocyclic, bicyclic or tricyclic partially saturated heterocycle; or a monocyclic, bicyclic or tricyclic aromatic heterocycle; each of the ring systems being substituted with formyl can be converted into compounds of

35

-31-


formula (I) wherein R^3 represents a monocyclic, bicyclic or tricyclic saturated carbocycle; a monocyclic, bicyclic or tricyclic partially saturated carbocycle; a monocyclic, bicyclic or tricyclic aromatic carbocycle; a monocyclic, bicyclic or tricyclic saturated heterocycle; a monocyclic, bicyclic or tricyclic partially saturated heterocycle; or a monocyclic, bicyclic or tricyclic aromatic heterocycle; each of the ring systems being substituted with $CH(=N-O-R^8)$ by reaction with NH_2OR^8 in the presence of a suitable base, such as for example sodium hydroxide and a suitable solvent, such as for example an alcohol, e.g. ethanol and the like. Compounds of formula (I) wherein R^3 represents a monocyclic, bicyclic or tricyclic saturated carbocycle; a monocyclic, bicyclic or tricyclic partially saturated carbocycle; a monocyclic, bicyclic or tricyclic aromatic carbocycle; a monocyclic, bicyclic or tricyclic saturated heterocycle; a monocyclic, bicyclic or tricyclic partially saturated heterocycle; or a monocyclic, bicyclic or tricyclic aromatic heterocycle; each of the ring systems being substituted with $CH(=N-O-R^8)$ can be converted into a compound of formula (I) wherein R^3 represents a monocyclic, bicyclic or tricyclic saturated carbocycle; a monocyclic, bicyclic or tricyclic partially saturated carbocycle; a monocyclic, bicyclic or tricyclic aromatic carbocycle; a monocyclic, bicyclic or tricyclic saturated heterocycle; a monocyclic, bicyclic or tricyclic partially saturated heterocycle; or a monocyclic, bicyclic or tricyclic aromatic heterocycle; each of the ring systems being substituted with CN by reaction with a carbodiimide in the presence of a suitable solvent, such as for example tetrahydrofuran.

Compounds of formula (I) wherein R^4 represents nitro, can be converted into a compound of formula (I) wherein R^4 is amino, in the presence of a suitable reducing agent, such as for example H_2 , in the presence of a suitable catalyst, such as for example Raney Nickel, and in the presence of a suitable solvent, such as for example an alcohol, e.g. methanol.


Compounds of formula (I) wherein R^1 is hydrogen, can be converted into a compound of formula (I) wherein R^1 is C_{1-6} alkyl, by reaction with a suitable alkylating agent, such as for example iodo- C_{1-6} alkyl, in the presence of a suitable base, such as for example sodium hydride, and a suitable solvent, such as for example tetrahydrofuran.

Compounds of formula (I-a) or (I-b) can be converted into a compound of formula (I) wherein R^{17} represents hydrogen, said compounds being represented by formula (I-c) and (I-d) by reaction with a suitable acid, such as for example trifluoroacetic acid.

-32-





- Compounds of formula (I-c) can be converted into a compound of formula (I) wherein R¹⁷ represents C₁₋₆alkyl optionally substituted with cyano or C₁₋₄alkyloxycarbonyl, said R¹⁷ being represented by formula R^{17c} and said compound being represented by formula (I-e), by reaction with an intermediate of formula R^{17c}-W₂, wherein W₂ represents a suitable leaving group, such as for example bromo, iodo and the like, in the presence of a suitable base, such as for example K₂CO₃, and a suitable solvent, such as for example acetonitrile.




- Compounds of formula (I-e) wherein R^{17c} represents C₁₋₆alkyl substituted with C₁₋₄alkyloxycarbonyl, said compounds being represented by formula (I-e-1), can be converted into a compound of formula (I) wherein R¹⁷ represents aminocarbonylC₁₋₆alkyl, said compound being represented by formula (I-f), by reaction with NH₃ in the presence of a suitable solvent, such as tetrahydrofuran or an alcohol, e.g. methanol.
- Compounds of formula (I-e-1) can also be converted into a compound of formula (I) wherein R¹⁷ represents hydroxyC₁₋₆alkyl, said compound being represented by formula (I-g), by reaction with NaBH₄ in the presence of a suitable solvent, such as for example tetrahydrofuran, an alcohol, e.g. ethanol.

-33-



Compounds of formula (I-f) can be converted into a compound of formula (I) wherein R¹⁷ represents cyanoC₁₋₆alkyl, said compound being represented by formula (I-h), by reaction with POCl₃ in the presence of a suitable solvent, such as for example

5 tetrahydrofuran.



Some of the compounds of formula (I) and some of the intermediates in the present invention may contain an asymmetric carbon atom. Pure stereochemically isomeric forms of said compounds and said intermediates can be obtained by the application of art-known procedures. For example, diastereoisomers can be separated by physical methods such as selective crystallization or chromatographic techniques, e.g. counter current distribution, liquid chromatography and the like methods. Enantiomers can be obtained from racemic mixtures by first converting said racemic mixtures with suitable resolving agents such as, for example, chiral acids, to mixtures of diastereomeric salts or compounds; then physically separating said mixtures of diastereomeric salts or


-34-

compounds by, for example, selective crystallization or chromatographic techniques, e.g. liquid chromatography and the like methods; and finally converting said separated diastereomeric salts or compounds into the corresponding enantiomers. Pure stereochemically isomeric forms may also be obtained from the pure stereochemically isomeric forms of the appropriate intermediates and starting materials, provided that the intervening reactions occur stereospecifically.

An alternative manner of separating the enantiomeric forms of the compounds of formula (I) and intermediates involves liquid chromatography, in particular liquid chromatography using a chiral stationary phase.

Some of the intermediates and starting materials are known compounds and may be commercially available or may be prepared according to art-known procedures or some of the compounds of formula (I) or the described intermediates may be prepared according to the procedures described in WO 99/50250, WO 00/27825 and WO03/016306.


Intermediates of formula (II-a) or (II-b) can be prepared by reacting an intermediate of formula (IV-a) or (IV-b) wherein W_1 is as defined above, with an intermediate of formula (III') in the presence of a suitable solvent, such as for example an alcohol, e.g. 2-butanol, a mixture of an alcohol and water, e.g. EtOH and water, or tetrahydrofuran, optionally in the presence of a suitable base, such as for example K_2CO_3 , $K^+ t\text{-}Bu^-$, collidine.



-35-

Intermediates of formula (II-a) or (II-b) wherein R^3 represents C_{2-6} alkenyl substituted with aminocarbonyl, can be converted into an intermediate of formula (II-a) or (II-b) wherein R^3 represents C_{2-6} alkenyl substituted with cyano, in the presence of $POCl_3$.



- 5 Intermediates of formula (II-a) and (II-b) wherein B represents a radical of formula (b) and wherein X_1 represents NR^5 , said intermediates being represented by formula (II-a-1) and (II-b-1), can be prepared by reacting an intermediate of formula (IV-a) or (IV-b) with an intermediate of formula (V) in the presence of a suitable solvent, such as for example tetrahydrofuran, t-amylOH or 1,2-dimethoxy ethane, and a suitable base, such as for example K_2CO_3 .




- 15 Intermediates of formula (II'-a) and (II'-b) wherein A represents a radical of formula (a) wherein R^1 represents hydrogen, said intermediates being represented by formula (II'-a-1) and (II'-b-1), can be prepared by reacting an intermediate of formula (VI-a) or (VI-b) with an intermediate of formula (VI') wherein W_{1a} represents a suitable leaving group, such as for example halogen, e.g. iodo, chloro and the like, in the presence of a suitable catalyst, such as for example $Pd(OAc)_2$, a suitable ligand, such as for example a mixture of (+) and (-) 2,2'-bis(diphenylphosphino)-1,1'-binaphthyl, a suitable base,

-36-


such as for example Cs_2CO_3 , and a suitable solvent, such as for example toluene or dioxane.



Intermediates of formula (IV-a) and (IV-b) can be prepared by reacting an intermediate of formula (VII) wherein W_1 is as defined hereinabove, with an intermediate of formula (VIII) wherein W_3 represents a suitable leaving group, such as for example halogen, e.g. chloro, iodo and the like, in the presence of a suitable solvent, such as for example *N,N*-dimethylformamide or acetonitrile, and a suitable base, such as for example K_2CO_3 .




Intermediates of formula (VI-a) and (VI-b) can be prepared by reacting an intermediate of formula (IX) with an intermediate of formula (VIII) in the presence of a suitable solvent, such as for example *N,N*-dimethylformamide or acetonitrile, and a suitable base, such as for example K_2CO_3 .




-37-

Intermediates of formula (V) can be prepared by reacting an intermediate of formula (X) with trifluoroacetic anhydride in the presence of a suitable solvent, such as for example 1,2-dimethoxy ethane.



- 5 Intermediates of formula (III) or (III') wherein A respectively B represents a radical of formula (b) wherein R^3 represents substituted C_{2-6} alkenyl, said R^3 being represented by C_{2-6} alkenyl- R^{3a} and said intermediates being represented by formula by formula (XI), can be prepared by reacting an intermediate of formula (XII) wherein W_4 represents a suitable leaving group, such as for example halogen, e.g. bromo and the like, with an
- 10 intermediate of formula (XIII) in the presence of a suitable catalyst, such as for example $Pd(OAc)_2$, a suitable ligand, such as for example tris(2-methylphenyl)phosphine, a suitable base, such as for example N,N -diethylethanamine, and a suitable solvent, such as for example MeCN.



- 15 Intermediates of formula (XI) wherein R^{3a} represents $-C(=O)-NH_2$, can be converted into an intermediate of formula (XI) wherein R^{3a} represents CN by reaction with $POCl_3$, optionally in the presence of dichloromethane.


- 20 Intermediates of formula (XI) wherein R^3 represents C_{2-6} alkenyl-CN, can be converted into an intermediate of formula (XI) wherein R^3 represents C_{2-6} alkyl-CN by reaction with H_2 , in the presence of a suitable catalyst, such as for example palladium on charcoal, and in the presence of a suitable solvent, such as for example an alcohol, e.g. ethanol.

-38-


Intermediates of formula (XI) wherein R^3 represents halo, can be converted into an intermediate of formula (XI) wherein R^3 represents $C(=O)-CH_3$, by reaction with n-butylvinyl ether, in the presence of a suitable catalyst, such as for example $Pd(OAc)_2$, a suitable ligand, such as for example 1,3-bis(diphenylphosphino)propane, a suitable base, such as for example K_2CO_3 , and a suitable solvent, such as for example N,N-dimethylformamide and H_2O .

Intermediates of formula (XI) wherein R^3 represents $C(=O)H$ respectively $C(=O)-CH_3$, can be converted into an intermediate of formula (XI) wherein R^3 represents $CH=CH-CN$ respectively $C(CH_3)=CH-CN$, by reaction with cyanomethylphosphonic acid diethylester in the presence of a suitable alcoholate, such as for example sodium methanolate, and a suitable solvent, such as for example an alcohol, e.g. methanol.

Intermediates of formula (XII) wherein W_4 represents bromo, said intermediates being represented by formula (XII-a), can be prepared by reacting an intermediate of formula (XIV) with Br_2 in the presence of a suitable acid, such as for example acetic acid; and a suitable solvent, such as an alcohol, e.g. methanol.




Intermediates of formula (III) or (III') wherein A respectively B represents a radical of formula (b) wherein X_1 represents NH , said intermediates being represented by formula (XV), can be prepared by reducing an intermediate of formula (XV') in the presence of a suitable reducing agent, such as for example Fe , in the presence of NH_4Cl , and a suitable solvent, such as for example alcohol/water mixture, e.g. $MeOH/H_2O$ (1:2).




Intermediates of formula (XV') wherein R^3 represents substituted C_{1-6} alkyloxy, said R^3 being represented by $R^{3a}-C_{1-6}$ alkyloxy, and said intermediates being represented by

-39-


formula (XV'-a), can be prepared by reacting an intermediate of formula (XVI) with an intermediate of formula (XVII) wherein W_5 represents a suitable leaving group, such as for example halogen, e.g. chloro and the like, in the presence of a suitable base, such as NaI, K_2CO_3 , and a suitable solvent, such as for example acetone.



Intermediates of formula (XVI) can be prepared by reacting an intermediate of formula (XVIII) with $NaNO_3$ in the presence of a suitable solvent, such as for example $MeSO_3H$.




- 10 Intermediates of formula (III) or (III') wherein A respectively B represents a radical of formula (b) wherein X_1 represents O, said intermediates being represented by formula (XIX), can be prepared by deprotecting an intermediate of formula (XX) wherein P represents a suitable protecting group, such as for example $-Si(CH_3)_2C(CH_3)_3$ or those protecting groups mentioned in Chapter 7 of 'Protective Groups in Organic Synthesis' by T Greene and P. Wuyts (John Wiley & Sons Inc., 1991), in the presence of 4-methylbenzenesulfonic acid and a suitable solvent, such as for example an alcohol, e.g. methanol.




- 20 Intermediates of formula (XX) wherein R^3 represents $CH=CH-CN$ respectively $C(CH_3)=CH-CN$, said intermediates being represented by formula (XX-a) respectively (XX-b), can be prepared by reacting an intermediate of formula (XXI-a) respectively (XXI-b) with cyanomethylphosphonic acid diethylester in the presence of a suitable

-40-

alcoholate, such as for example sodium methanolate, and a suitable solvent, such as for example tetrahydrofuran.



- 5 Intermediates of formula (XXI-a) respectively (XXI-b) can be prepared by reacting an intermediate of formula (XXII) with an intermediate of formula (XXIII) wherein W_6 represents a suitable leaving group, such as for example halogen, e.g. chloro and the like, and P is as defined hereinabove, in the presence of a suitable solvent, such as for example *N,N*-dimethylformamide, and a suitable base, such as for example diisopropylethanamine.



10

It will be appreciated by those skilled in the art that in the processes described above the functional groups of intermediate compounds may need to be blocked by protecting groups.

-41-

Functional groups which it is desirable to protect include hydroxy, amino and carboxylic acid. Suitable protecting groups for hydroxy include trialkylsilyl groups (e.g. *tert*-butyldimethylsilyl, *tert*-butyldiphenylsilyl or trimethylsilyl), benzyl and tetrahydropyranyl. Suitable protecting groups for amino include *tert*-butoxycarbonyl or benzyloxycarbonyl. Suitable protecting groups for carboxylic acid include C₁₋₆alkyl or benzyl esters.

The protection and deprotection of functional groups may take place before or after a reaction step.

The use of protecting groups is fully described in 'Protective Groups in Organic Chemistry', edited by J W F McOmie, Plenum Press (1973), and 'Protective Groups in Organic Synthesis' 2nd edition, T W Greene & P G M Wutz, Wiley Interscience (1991).

The compounds of formula (I) or any subgroup thereof show antiretroviral properties (reverse transcriptase inhibiting properties), in particular against Human Immunodeficiency Virus (HIV), which is the aetiological agent of Acquired Immune Deficiency Syndrome (AIDS) in humans. The HIV virus preferentially infects human T-4 cells and destroys them or changes their normal function, particularly the coordination of the immune system. As a result, an infected patient has an ever decreasing number of T-4 cells, which moreover behave abnormally. Hence, the immunological defense system is unable to combat infections and neoplasms and the HIV infected subject usually dies by opportunistic infections such as pneumonia, or by cancers. Other conditions associated with HIV infection include thrombocytopaenia, Kaposi's sarcoma and infection of the central nervous system characterized by progressive demyelination, resulting in dementia and symptoms such as, progressive dysarthria, ataxia and disorientation. HIV infection further has also been associated with peripheral neuropathy, progressive generalized lymphadenopathy (PGL) and AIDS-related complex (ARC).

The present compounds also show activity against drug or multidrug resistant HIV strains, in particular drug or multidrug resistant HIV-1 strains, more in particular the present compounds show activity against HIV strains, especially HIV-1 strains, that have acquired resistance to one or more art-known non-nucleoside reverse transcriptase inhibitors. Art-known non-nucleoside reverse transcriptase inhibitors are those non-nucleoside reverse transcriptase inhibitors other than the present compounds and in

-42-

particular commercial non-nucleoside reverse transcriptase inhibitors. The present compounds also have little or no binding affinity to human α -1 acid glycoprotein; human α -1 acid glycoprotein does not or only weakly affect the anti HIV activity of the present compounds.

5

Due to their antiretroviral properties, particularly their anti-HIV properties, especially their anti-HIV-1-activity, the compounds of formula (I) or any subgroup thereof, their *N*-oxides, pharmaceutically acceptable addition salts, quaternary amines or stereochemically isomeric forms thereof, are useful in the treatment of individuals infected by HIV and for the prophylaxis of these infections. In general, the compounds of the present invention may be useful in the treatment of warm-blooded mammals infected with viruses whose existence is mediated by, or depends upon, the enzyme reverse transcriptase. Conditions which may be prevented or treated with the compounds of the present invention, especially conditions associated with HIV and other pathogenic retroviruses, include AIDS, AIDS-related complex (ARC), progressive generalized lymphadenopathy (PGL), as well as chronic Central Nervous System diseases caused by retroviruses, such as, for example HIV mediated dementia and multiple sclerosis.

20 The compounds of the present invention or any subgroup thereof may therefore be used as medicines against above-mentioned conditions. Said use as a medicine or method of treatment comprises the administration to HIV-infected subjects of an amount effective to combat the conditions associated with HIV and other pathogenic retroviruses, especially HIV-1. In particular, the compounds of formula (I) or any subgroup thereof, their *N*-oxides, pharmaceutically acceptable addition salts, quaternary amines or stereochemically isomeric forms thereof, may be used for the manufacture of a medicament for the treatment or the prevention of HIV infections.

30 In view of the utility of the compounds of formula (I) or any subgroup thereof, there is provided a method of treating warm-blooded mammals, including humans, suffering from or a method of preventing warm-blooded mammals, including humans, to suffer from viral infections, especially HIV infections. Said method comprises the administration, preferably oral administration, of an effective amount of a compound of formula (I) or any subgroup thereof, a *N*-oxide form, a pharmaceutically acceptable addition salt, a quaternary amine or a possible stereoisomeric form thereof, to warm-blooded mammals, including humans.

35

-43-

5 The present invention also provides compositions for treating viral infections comprising a therapeutically effective amount of a compound of formula (I) or any subgroup thereof, a *N*-oxide, pharmaceutically acceptable addition salt, quaternary amine or stereochemically isomeric form thereof, and a pharmaceutically acceptable carrier or diluent.

10 The compounds of the present invention or any subgroup thereof may be formulated into various pharmaceutical forms for administration purposes. As appropriate compositions there may be cited all compositions usually employed for systemically administering drugs. To prepare the pharmaceutical compositions of this invention, an effective amount of the particular compound, optionally in addition salt form, as the active ingredient is combined in intimate admixture with a pharmaceutically acceptable carrier, which carrier may take a wide variety of forms depending on the form of preparation desired for administration. These pharmaceutical compositions are
15 desirable in unitary dosage form suitable, particularly, for administration orally, rectally, percutaneously, or by parenteral injection. For example, in preparing the compositions in oral dosage form, any of the usual pharmaceutical media may be employed such as, for example, water, glycols, oils, alcohols and the like in the case of oral liquid preparations such as suspensions, syrups, elixirs, emulsions and solutions; or
20 solid carriers such as starches, sugars, kaolin, diluents, lubricants, binders, disintegrating agents and the like in the case of powders, pills, capsules, and tablets. Because of their ease in administration, tablets and capsules represent the most advantageous oral dosage unit forms, in which case solid pharmaceutical carriers are obviously employed. For parenteral compositions, the carrier will usually comprise
25 sterile water, at least in large part, though other ingredients, for example, to aid solubility, may be included. Injectable solutions, for example, may be prepared in which the carrier comprises saline solution, glucose solution or a mixture of saline and glucose solution. Injectable suspensions may also be prepared in which case appropriate liquid carriers, suspending agents and the like may be employed. Also
30 included are solid form preparations which are intended to be converted, shortly before use, to liquid form preparations. In the compositions suitable for percutaneous administration, the carrier optionally comprises a penetration enhancing agent and/or a suitable wetting agent, optionally combined with suitable additives of any nature in minor proportions, which additives do not introduce a significant deleterious effect on
35 the skin. Said additives may facilitate the administration to the skin and/or may be helpful for preparing the desired compositions. These compositions may be administered in various ways, e.g., as a transdermal patch, as a spot-on, as an ointment.

-44-

The compounds of the present invention may also be administered via inhalation or insufflation by means of methods and formulations employed in the art for administration via this way. Thus, in general the compounds of the present invention may be administered to the lungs in the form of a solution, a suspension or a dry powder. Any system developed for the delivery of solutions, suspensions or dry powders via oral or nasal inhalation or insufflation are suitable for the administration of the present compounds.

To aid solubility of the compounds of formula (I) or any subgroup thereof, suitable ingredients, e.g. cyclodextrins, may be included in the compositions. Appropriate cyclodextrins are α -, β -, γ -cyclodextrins or ethers and mixed ethers thereof wherein one or more of the hydroxy groups of the anhydroglucose units of the cyclodextrin are substituted with C₁₋₆alkyl, particularly methyl, ethyl or isopropyl, e.g. randomly methylated β -CD; hydroxyC₁₋₆alkyl, particularly hydroxyethyl, hydroxy-propyl or hydroxybutyl; carboxyC₁₋₆alkyl, particularly carboxymethyl or carboxy-ethyl; C₁₋₆alkylcarbonyl, particularly acetyl. Especially noteworthy as complexants and/or solubilizers are β -CD, randomly methylated β -CD, 2,6-dimethyl- β -CD, 2-hydroxyethyl- β -CD, 2-hydroxyethyl- β -CD, 2-hydroxypropyl- β -CD and (2-carboxymethoxy)propyl- β -CD, and in particular 2-hydroxypropyl- β -CD (2-HP- β -CD).

The term mixed ether denotes cyclodextrin derivatives wherein at least two cyclodextrin hydroxy groups are etherified with different groups such as, for example, hydroxy-propyl and hydroxyethyl.

The average molar substitution (M.S.) is used as a measure of the average number of moles of alkoxy units per mole of anhydroglucose. The average substitution degree (D.S.) refers to the average number of substituted hydroxyls per anhydroglucose unit. The M.S. and D.S. value can be determined by various analytical techniques such as nuclear magnetic resonance (NMR), mass spectrometry (MS) and infrared spectroscopy (IR). Depending on the technique used, slightly different values may be obtained for one given cyclodextrin derivative. Preferably, as measured by mass spectrometry, the M.S. ranges from 0.125 to 10 and the D.S. ranges from 0.125 to 3.

Other suitable compositions for oral or rectal administration comprise particles consisting of a solid dispersion comprising a compound of formula (I) or any subgroup

-45-

thereof and one or more appropriate pharmaceutically acceptable water-soluble polymers.

5 The term "a solid dispersion" used hereinafter defines a system in a solid state (as opposed to a liquid or gaseous state) comprising at least two components, in casu the compound of formula (I) or any subgroup thereof and the water-soluble polymer, wherein one component is dispersed more or less evenly throughout the other component or components (in case additional pharmaceutically acceptable formulating agents, generally known in the art, are included, such as plasticizers, preservatives and the like). When said dispersion of the components is such that the system is chemically and physically uniform or homogenous throughout or consists of one phase as defined in thermo-dynamics, such a solid dispersion will be called "a solid solution". Solid solutions are preferred physical systems because the components therein are usually readily bioavailable to the organisms to which they are administered. This advantage can probably be explained by the ease with which said solid solutions can form liquid solutions when contacted with a liquid medium such as the gastro-intestinal juices. The ease of dissolution may be attributed at least in part to the fact that the energy required for dissolution of the components from a solid solution is less than that required for the dissolution of components from a crystalline or microcrystalline solid phase.

20 The term "a solid dispersion" also comprises dispersions which are less homogenous throughout than solid solutions. Such dispersions are not chemically and physically uniform throughout or comprise more than one phase. For example, the term "a solid dispersion" also relates to a system having domains or small regions wherein amorphous, microcrystalline or crystalline compound of formula (I) or any subgroup thereof, or amorphous, microcrystalline or crystalline water-soluble polymer, or both, are dispersed more or less evenly in another phase comprising water-soluble polymer, or compound of formula (I) or any subgroup thereof, or a solid solution comprising compound of formula (I) or any subgroup thereof and water-soluble polymer. Said domains are regions within the solid dispersion distinctively marked by some physical feature, small in size, and evenly and randomly distributed throughout the solid dispersion.

35 Various techniques exist for preparing solid dispersions including melt-extrusion, spray-drying and solution-evaporation.

The solution-evaporation process comprises the following steps :

-46-

- a) dissolving the compound of formula (I) or any subgroup thereof and the water-soluble polymer in an appropriate solvent, optionally at elevated temperatures;
b) heating the solution resulting under point a), optionally under vacuum, until the solvent is evaporated. The solution may also be poured onto a large surface so as to
5 form a thin film, and evaporating the solvent therefrom.

In the spray-drying technique, the two components are also dissolved in an appropriate solvent and the resulting solution is then sprayed through the nozzle of a spray dryer followed by evaporating the solvent from the resulting droplets at elevated
10 temperatures.

The preferred technique for preparing solid dispersions is the melt-extrusion process comprising the following steps :

- a) mixing a compound of formula (I) or any subgroup thereof and an appropriate
15 water-soluble polymer,
b) optionally blending additives with the thus obtained mixture,
c) heating and compounding the thus obtained blend until one obtains a
homogenous melt,
d) forcing the thus obtained melt through one or more nozzles; and
20 e) cooling the melt till it solidifies.

The terms "melt" and "melting" should be interpreted broadly. These terms not only mean the alteration from a solid state to a liquid state, but can also refer to a transition to a glassy state or a rubbery state, and in which it is possible for one component of the
25 mixture to get embedded more or less homogeneously into the other. In particular cases, one component will melt and the other component(s) will dissolve in the melt thus forming a solution, which upon cooling may form a solid solution having advantageous dissolution properties.

30 After preparing the solid dispersions as described hereinabove, the obtained products can be optionally milled and sieved.

The solid dispersion product may be milled or ground to particles having a particle size of less than 600 μm , preferably less than 400 μm and most preferably less than 125 μm .
35

The particles prepared as described hereinabove can then be formulated by conventional techniques into pharmaceutical dosage forms such as tablets and capsules.

-47-

It will be appreciated that a person of skill in the art will be able to optimize the parameters of the solid dispersion preparation techniques described above, such as the most appropriate solvent, the working temperature, the kind of apparatus being used,
5 the rate of spray-drying, the throughput rate in the melt-extruder

The water-soluble polymers in the particles are polymers that have an apparent viscosity, when dissolved at 20°C in an aqueous solution at 2 % (w/v), of 1 to 5000 mPa.s more preferably of 1 to 700 mPa.s, and most preferred of 1 to 100 mPa.s. For
10 example, suitable water-soluble polymers include alkylcelluloses, hydroxyalkyl-celluloses, hydroxyalkyl alkylcelluloses, carboxyalkylcelluloses, alkali metal salts of carboxyalkylcelluloses, carboxyalkylalkylcelluloses, carboxyalkylcellulose esters, starches, pectines, chitin derivatives, di-, oligo- and polysaccharides such as trehalose, alginic acid or alkali metal and ammonium salts thereof, carrageenans, galactomannans,
15 tragacanth, agar-agar, gummi arabicum, guar gummi and xanthan gummi, polyacrylic acids and the salts thereof, polymethacrylic acids and the salts thereof, methacrylate copolymers, polyvinylalcohol, polyvinylpyrrolidone, copolymers of polyvinylpyrrolidone with vinyl acetate, combinations of polyvinylalcohol and polyvinylpyrrolidone, polyalkylene oxides and copolymers of ethylene oxide and
20 propylene oxide. Preferred water-soluble polymers are hydroxypropyl methylcelluloses.

Also one or more cyclodextrins can be used as water soluble polymer in the preparation of the above-mentioned particles as is disclosed in WO 97/18839. Said cyclodextrins
25 include the pharmaceutically acceptable unsubstituted and substituted cyclodextrins known in the art, more particularly α , β or γ cyclodextrins or the pharmaceutically acceptable derivatives thereof.

Substituted cyclodextrins which can be used to prepare the above described particles
30 include polyethers described in U.S. Patent 3,459,731. Further substituted cyclodextrins are ethers wherein the hydrogen of one or more cyclodextrin hydroxy groups is replaced by C₁₋₆alkyl, hydroxyC₁₋₆alkyl, carboxy-C₁₋₆alkyl or C₁₋₆alkyloxycarbonylC₁₋₆alkyl or mixed ethers thereof. In particular such substituted cyclodextrins are ethers wherein the hydrogen of one or more cyclodextrin hydroxy
35 groups is replaced by C₁₋₃alkyl, hydroxyC₂₋₄alkyl or carboxyC₁₋₂alkyl or more in particular by methyl, ethyl, hydroxyethyl, hydroxypropyl, hydroxybutyl, carboxy-methyl or carboxyethyl.

-48-

Of particular utility are the β -cyclodextrin ethers, e.g. dimethyl- β -cyclodextrin as described in Drugs of the Future, Vol. 9, No. 8, p. 577-578 by M. Nogradi (1984) and polyethers, e.g. hydroxypropyl β -cyclodextrin and hydroxyethyl β -cyclodextrin, being
5 examples. Such an alkyl ether may be a methyl ether with a degree of substitution of about 0.125 to 3, e.g. about 0.3 to 2. Such a hydroxypropyl cyclodextrin may for example be formed from the reaction between β -cyclodextrin an propylene oxide and may have a MS value of about 0.125 to 10, e.g. about 0.3 to 3.

10 Another type of substituted cyclodextrins is sulfobutylcyclodextrines.

The ratio of the compound of formula (I) or any subgroup thereof over the water soluble polymer may vary widely. For example ratios of 1/100 to 100/1 may be applied. Interesting ratios of the compound of formula (I) or any subgroup thereof over
15 cyclodextrin range from about 1/10 to 10/1. More interesting ratios range from about 1/5 to 5/1.

It may further be convenient to formulate the compounds of formula (I) or any subgroup thereof in the form of nanoparticles which have a surface modifier adsorbed
20 on the surface thereof in an amount sufficient to maintain an effective average particle size of less than 1000 nm. Useful surface modifiers are believed to include those which physically adhere to the surface of the compound of formula (I) but do not chemically bond to said compound.

25 Suitable surface modifiers can preferably be selected from known organic and inorganic pharmaceutical excipients. Such excipients include various polymers, low molecular weight oligomers, natural products and surfactants. Preferred surface modifiers include nonionic and anionic surfactants.

30 Yet another interesting way of formulating the compounds of formula (I) or any subgroup thereof involves a pharmaceutical composition whereby the compounds of formula (I) or any subgroup thereof are incorporated in hydrophilic polymers and applying this mixture as a coat film over many small beads, thus yielding a composition which can conveniently be manufactured and which is suitable for preparing
35 pharmaceutical dosage forms for oral administration.

-49-

Said beads comprise a central, rounded or spherical core, a coating film of a hydrophilic polymer and a compound of formula (I) or any subgroup thereof and optionally a seal-coating layer.

- 5 Materials suitable for use as cores in the beads are manifold, provided that said materials are pharmaceutically acceptable and have appropriate dimensions and firmness. Examples of such materials are polymers, inorganic substances, organic substances, and saccharides and derivatives thereof.
- 10 It is especially advantageous to formulate the aforementioned pharmaceutical compositions in unit dosage form for ease of administration and uniformity of dosage. Unit dosage form as used herein refers to physically discrete units suitable as unitary dosages, each unit containing a predetermined quantity of active ingredient calculated to produce the desired therapeutic effect in association with the required
- 15 pharmaceutical carrier. Examples of such unit dosage forms are tablets (including scored or coated tablets), capsules, pills, powder packets, wafers, suppositories, injectable solutions or suspensions and the like, and segregated multiples thereof.

Those of skill in the treatment of HIV-infection could determine the effective daily

20 amount from the test results presented here. In general it is contemplated that an effective daily amount would be from 0.01 mg/kg to 50 mg/kg body weight, more preferably from 0.1 mg/kg to 10 mg/kg body weight. It may be appropriate to administer the required dose as two, three, four or more sub-doses at appropriate intervals throughout the day. Said sub-doses may be formulated as unit dosage forms,

25 for example, containing 1 to 1000 mg, and in particular 5 to 200 mg of active ingredient per unit dosage form.

The exact dosage and frequency of administration depends on the particular compound of formula (I) or any subgroup thereof, used, the particular condition being treated, the

30 severity of the condition being treated, the age, weight and general physical condition of the particular patient as well as other medication the individual may be taking, as is well known to those skilled in the art. Furthermore, it is evident that said effective daily amount may be lowered or increased depending on the response of the treated subject and/or depending on the evaluation of the physician prescribing the compounds

35 of the instant invention. The effective daily amount ranges mentioned hereinabove are therefore only guidelines and are not intended to limit the scope or use of the invention to any extent.

-50-

The present compounds of formula (I) or any subgroup thereof can be used alone or in combination with other therapeutic agents, such as anti-virals, antibiotics, immunomodulators or vaccines for the treatment of viral infections. They may also be used alone or in combination with other prophylactic agents for the prevention of viral infections. The present compounds may be used in vaccines and methods for protecting individuals against viral infections over an extended period of time. The compounds may be employed in such vaccines either alone or together with other compounds of this invention or together with other anti-viral agents in a manner consistent with the conventional utilization of reverse transcriptase inhibitors in vaccines. Thus, the present compounds may be combined with pharmaceutically acceptable adjuvants conventionally employed in vaccines and administered in prophylactically effective amounts to protect individuals over an extended period of time against HIV infection.

Also, the combination of an antiretroviral compound and a compound of formula (I) or any subgroup thereof can be used as a medicine. Thus, the present invention also relates to a product containing (a) a compound of formula (I) or any subgroup thereof, and (b) another antiretroviral compound, as a combined preparation for simultaneous, separate or sequential use in anti-HIV treatment. The different drugs may be combined in a single preparation together with pharmaceutically acceptable carriers. Said other antiretroviral compounds may be known antiretroviral compounds such as suramine, pentamidine, thymopentin, castanospermine, dextran (dextran sulfate), foscarnet-sodium (trisodium phosphono formate); nucleoside reverse transcriptase inhibitors, e.g. zidovudine (3'-azido-3'-deoxythymidine, AZT), didanosine (2',3'-dideoxyinosine; ddI), zalcitabine (dideoxycytidine, ddC) or lamivudine (2'-3'-dideoxy-3'-thiacytidine, 3TC), stavudine (2',3'-didehydro-3'-deoxythymidine, d4T), abacavir and the like; non-nucleoside reverse transcriptase inhibitors such as nevirapine (11-cyclopropyl-5,11-dihydro-4-methyl-6H-dipyrido-[3,2-b : 2',3'-e][1,4]diazepin-6-one), efavirenz, delavirdine, TMC-120, TMC-125, 4-[[4-[[4-(2-cyanoethenyl)-2,6-dimethylphenyl]-amino]-2-pyrimidinyl]amino]benzonitrile, and the like; phosphonate reverse transcriptase inhibitors, e.g. tenofovir and the like; compounds of the TIBO (tetrahydro-imidazo[4,5,1-jk][1,4]-benzodiazepine-2(1H)-one and thione)-type e.g. (S)-8-chloro-4,5,6,7-tetrahydro-5-methyl-6-(3-methyl-2-butenyl)imidazo-[4,5,1-jk][1,4]benzodiazepine-2(1H)-thione; compounds of the α -APA (α -anilino phenyl acetamide) type e.g. α -[(2-nitrophenyl)amino]-2,6-dichlorobenzene-acetamide and the like; inhibitors of trans-activating proteins, such as

-51-

TAT-inhibitors, e.g. RO-5-3335, or REV inhibitors, and the like; protease inhibitors e.g. indinavir, ritonavir, saquinavir, lopinavir (ABT-378), nelfinavir, amprenavir, TMC-126, BMS-232632, VX-175 and the like; fusion inhibitors, e.g. T-20, T-1249 and the like; CXCR4 receptor antagonists, e.g. AMD-3100 and the like; inhibitors of the viral integrase; nucleotide-like reverse transcriptase inhibitors, e.g. tenofovir and the like; ribonucleotide reductase inhibitors, e.g. hydroxyurea and the like.

By administering the compounds of the present invention with other anti-viral agents which target different events in the viral life cycle, the therapeutic effect of these compounds can be potentiated. Combination therapies as described above exert a synergistic effect in inhibiting HIV replication because each component of the combination acts on a different site of HIV replication. The use of such combinations may reduce the dosage of a given conventional anti-retroviral agent which would be required for a desired therapeutic or prophylactic effect as compared to when that agent is administered as a monotherapy. These combinations may reduce or eliminate the side effects of conventional single anti-retroviral therapy while not interfering with the anti-viral activity of the agents. These combinations reduce potential of resistance to single agent therapies, while minimizing any associated toxicity. These combinations may also increase the efficacy of the conventional agent without increasing the associated toxicity.

The compounds of the present invention may also be administered in combination with immunomodulating agents, e.g. levamisole, bropirimine, anti-human alpha interferon antibody, interferon alpha, interleukin 2, methionine enkephalin, diethyldithiocarbamate, tumor necrosis factor, naltrexone and the like; antibiotics, e.g. pentamidine isethionate and the like; cholinergic agents, e.g. tacrine, rivastigmine, donepezil, galantamine and the like; NMDA channel blockers, e.g. memantine to prevent or combat infection and diseases or symptoms of diseases associated with HIV infections, such as AIDS and ARC, e.g. dementia. A compound of formula (I) or any subgroup thereof can also be combined with another compound of formula (I) or any subgroup thereof.

Although the present invention focuses on the use of the present compounds for preventing or treating HIV infections, the present compounds may also be used as inhibitory agents for other viruses which depend on similar reverse transcriptases for obligatory events in their life cycle.

-52-

The following examples illustrate the present invention.


Experimental part

Hereinafter, "DMF" is defined as *N,N*-dimethylformamide, "DME" is defined as 1,2-dimethoxyethane, "THF" is defined as tetrahydrofuran, "BINAP" is defined as a mixture of (+) and (-) [1,1'-binaphthalene]-2,2'-diylbis[diphenylphosphine], "TFA" is defined as trifluoroacetic acid, "TFAA" is defined as trifluoroacetic anhydride, "DCE" is defined as 1,2-dichloro ethane and "DIPEA" is defined as diisopropylethanamine.

10 A. Preparation of the intermediate compounds


Example A1

a) Preparation of intermediate 1



2-Methoxy-6-methylbenzenamine (12.12 g) was dissolved in 100 ml of MeOH and 10 ml of HOAc. Temperature was lowered to 0°C. A solution of 14.12 g Br₂ in 50 ml of MeOH was added with careful temperature-control. After addition, the solvents were
15 evaporated and the residue was dissolved in 300 ml of diethyl ether and 100 ml of 2M NaOH. The layers were separated. The ether-layer was dried (NaCl (saturated) and Na₂SO₄), filtered and concentrated. Yield: 18.68 g of intermediate 1 (4-bromo-2-methoxy-6-methylbenzenamine) (98%) .

b) Preparation of intermediate 2




0.1 eq of Pd(OAc)₂ (1.23 g), 0.3 eq of tris(2-methylphenyl)phosphine (5.0 g), 1.5 eq of Et₃N (8.31 g), 4-bromo-2-methoxy-6-methylbenzenamine (intermediate 1) (11.83 g)
20 and 2-propenamide (1.5 eq., 5.84 g) were brought in 150 ml of MeCN and N₂ was bubbled through the suspension for at least 20 minutes. Then a cooler was mounted strictly under nitrogen atmosphere. The reaction was heated at 70°C overnight. The reaction mixture was allowed to cool to 20°C and was diluted with 700 ml of CH₂Cl₂
25 and washed with saturated aqueous NaHCO₃ (2x 100 ml) and dried with brine and

-53-

Na₂SO₄. The residue was triturated in *i*Pr₂O, filtered off and air-dried. Yield: 11.65 g of intermediate 2 (99%).


c) Preparation of intermediate 3



Intermediate 2 (11.65 g) was dissolved in 50 ml of POCl₃ at 20°C. The reaction mixture was stirred at 20°C and checked by TLC. The reaction mixture was added dropwise to 500 ml of diisopropyl ether. The precipitate was filtered off, the residue was added to 350 ml of EtOAc and 250 ml of NaHCO₃ (saturated). The layers were separated. The water-layer was washed with 150 ml of EtOAc once. The combined EtOAc-layers were dried (NaCl (saturated) and Na₂SO₄), filtered and concentrated. Yield: 9.55 g of intermediate 3 (85%).

10 Example A2


Preparation of intermediate 4




0.1 eq (335 mg) of Pd(OAc)₂, 0.2 eq (908 mg) of tris(2-methylphenyl)phosphine, 1.5 eq (3.11 ml) of Et₃N, 4-bromo-2,6-dimethylphenol (3.00 g) and *N,N*-dimethyl-2-propenamide (1.5 eq., 2.31 ml) were brought in 100 ml of MeCN and N₂ was bubbled through the suspension for at least 20 minutes. Then a cooler was mounted strictly under nitrogen atmosphere. The reaction was heated at 70°C overnight. The reaction mixture was allowed to cool to 20°C and was diluted with 500 ml of CH₂Cl₂ and washed with saturated aqueous NaHCO₃ (2x100 ml) and dried with brine and Na₂SO₄. The residue was sonicated in *i*Pr₂O and filtered off. Yield: 2.26 g of intermediate 4 (69%).

20 Example A3

a) Preparation of intermediate 6




-54-



Intermediate 5 (prepared according to A1b) (6.81 g) was dissolved in 26 ml of POCl_3 at 20°C . The reaction mixture was stirred at 20°C and checked by TLC. The reaction mixture was added dropwise to 500 ml of diisopropyl ether. The precipitate was filtered off, the residue was added to 250 ml of EtOAc and 150 ml of NaHCO₃ (saturated) (aqueous). The layers were separated. The water-layer was washed with 100 ml of EtOAc once. The combined EtOAc-layers were dried (NaCl (saturated) and Na₂SO₄), filtered and concentrated. Yield: 5.37 g of intermediate 6 (87%).


b) Preparation of intermediate 7



Hydrogenation at $P = P^0$ using 0.05 eq (0.88 g) of 5% Pd/C in 200 ml of EtOH with 1.40 gram of intermediate 6. After 4 hours, the Pd/C was filtered off and the filtrate was evaporated and stripped with 50 ml of EtOAc and with 50 ml of CH₂Cl₂. Yield : 1.35 g of intermediate 7 (93%).

Example A4


Preparation of intermediate 8



Intermediate 3 (3.81 g) was suspended in 100 ml of 1,2-dimethoxyethane (DME) at 0°C and stirred vigorously. At 0°C , a solution of TFAA (2.0 eq., 8.48 g) in DME was added dropwise. After vigorously stirring for 30 minutes at 0°C , the reaction mixture was stirred vigorously at 20°C and checked by TLC. The reaction was quenched by adding 400 ml of NaHCO₃ (saturated) (aqueous). EtOAc (200 ml) was added to the quenched reaction mixture. The layers were separated. The water-layer was washed with EtOAc (100 ml) once. The combined EtOAc-layers were dried (NaCl (saturated) and Na₂SO₄), filtered and concentrated. Yield : 5.77 g of intermediate 8 (98%).

Example A5


a) Preparation of intermediate 9



-55-


3,5-Dimethylphenol (50 g) was dissolved in 200 ml of MeSO_3H and cooled to 0°C . 1 eq (34.8 g) of NaNO_3 was added portion-wise at 0°C . After 18 hours, the reaction mixture was poured into 4 litre of ice- H_2O under vigorous stirring. The water was decanted. The residue was dissolved in 400 ml of EtOAc. The EtOAc extract was washed with saturated aqueous NaHCO_3 (2x200 ml), dried using brine and Na_2SO_4 , and purified using a mixture of EtOAc and n-heptane (4:1) on silicagel. Yield: 7.86 g of intermediate 9 (11%).

b) Preparation of intermediate 10



Intermediate 9 (1.40 g) was dissolved in 40 ml of acetone. Subsequently, K_2CO_3 (2.0 eq., 2.31 g) and NaI (0.1 eq., 126 mg) were added, followed by ClCH_2CN (1.5 eq., 0.95 g). After completion of the reaction (GC), the reaction-mixture was filtered and concentrated. The residue was dissolved in 100 ml of EtOAc. The EtOAc extract was washed with saturated aqueous NaHCO_3 (2x200 ml), dried using brine and Na_2SO_4 . Yield: 1.91 g of intermediate 10 (99%).

c) Preparation of intermediate 11




5 eq. (1.30 g) of NH_4Cl was dissolved in 20 ml of H_2O and 3 eq (0.81 g) of Fe was added. Intermediate 10 (1.00 g) was dissolved in 40 ml of MeOH . The solution was added to the aqueous solution. The reaction was stirred at 50°C and checked by TLC. The reaction-mixture was filtered hot. The filtrate was poured in 200 ml of EtOAc. The EtOAc extract was washed with saturated aqueous NaHCO_3 (2x200 ml), dried using brine and Na_2SO_4 . Yield : 0.92 g of intermediate 11 (100%).

-56-

Example A6-1


a) Preparation of intermediate 12




4-Hydroxy-3,5-dimethoxybenzaldehyde (2.00 g) and DIPEA (2.0 eq., 2.84 g) was dissolved in 25 ml of DMF. A solution of chloro(1,1-dimethylethyl)dimethylsilane (1.1 eq., 1.82 g) in 10 ml of DMF was added dropwise. After 18 hours, the reaction mixture was poured in 150 ml of water, followed by extraction with ethyl ether (2x100 ml). The organic layer was dried (NaCl (saturated) and Na₂SO₄), filtered and concentrated. Yield: 2.91 g of intermediate 12 (97%):

b) Preparation of intermediate

13 and 14



Intermediate 13




Intermediate 14

NaOMe (2.0 eq., 0.73 g) was suspended in 20 ml of THF. A solution of intermediate 12 (2.00 g) in 10 ml of THF was added dropwise. After stirring for 15 minutes, a solution of cyanomethylphosphonic acid diethyl ester (1.0 eq., 1.20 g) was added dropwise. After stirring for 18 hours, the reaction was quenched by adding 0.5 M HCl until the pH was below 1. The quenched reaction mixture was extracted with 150 ml of EtOAc. The layers were separated. The water-layer was washed with 50 ml of EtOAc once. The combined EtOAc-layers were dried (NaCl (saturated) and Na₂SO₄), filtered and concentrated. The residue was dissolved in ethyl ether (50 ml). By adding heptane, the main product from the reaction (the trans-isomer, intermediate 13) precipitated. The mother liquid contained both the cis- and the trans-isomer. Yield : 0.58 g of intermediate 13 (27%).

-57-


c) Preparation of intermediate 15



Intermediate 13 (580 mg) was dissolved in 15 ml of MeOH. p-Toluenesulfonic acid (0.05 eq., 20 mg) was added. After 40 hours at 60°C, the deprotection was complete. The reaction-mixture was poured in 50 ml of water and extracted with 150 ml of EtOAc. The layers were separated. The water-layer was washed with 50 ml of EtOAc once. The combined EtOAc-layers were washed with 50 ml of NaHCO₃ (saturated) (aqueous), dried (NaCl (saturated) and Na₂SO₄), filtered and concentrated. Yield : 0.33 g of intermediate 15 (89%).

Example A6-2


a) Preparation of intermediate 41




- 10 4-Bromo-2,6-dimethylaniline (4.0 gram), *n*-butyl vinyl ether (3.0 eq., 6.0 gram), K₂CO₃ (1.2 eq., 3.32 gram), Pd(OAc)₂ (0.03 eq., 135 mg) and 1,3-bis(diphenylphosphino)propane (0.66 eq., 0.54 gram) were dissolved in 25 mL of DMF and 3 mL of H₂O and N₂ was bubbled through the suspension for at least 20 minutes. After 24 hours at 80°C, the same amounts of palladium and 1,3-
- 15 bis(diphenylphosphino)propane were added. Subsequently, the reaction mixture was kept at 80°C for 16 hours. The reaction mixture was dissolved in 150 ml of Et₂O and 50 ml of 1 N HCl was added. After stirring for 1 hour, the acid aqueous layer was basified with 2 N NaOH. The layers were separated. The water-layer was washed with 50 ml of Et₂O once more. The combined Et₂O-layers were dried (NaCl (saturated) and
- 20 Na₂SO₄), filtered and concentrated. Yield :2.72 gr of intermediate 41.

-58-

b) Preparation of intermediate 42 and 43



Intermediate 42




(E)

Intermediate 43


NaOMe (3.0 eq., 0.33 g) was dissolved in 20 ml of MeOH. Intermediate 41 (0.33 g) was added in one portion. After stirring for 15 minutes, cyanomethyl phosphonic acid diethyl ester (1.5 eq., 0.53 g) was added dropwise. After stirring for 3 days at 20°C, another portion of cyanomethyl phosphonic acid diethyl ester (1.5 eq., 0.53 g) was added dropwise. After stirring for 4 days at 20°C, another portion of cyanomethylphosphonic acid diethyl ester (1.5 eq., 0.53 g) was added dropwise, followed by NaOMe (1.5 eq., 0.17 g). After total 7 days, the reaction mixture was concentrated. The residue was dissolved in 150 ml of EtOAc and 100 ml of aqueous NaHCO₃ (saturated). The layers were separated. The aqueous layer was washed with 50 ml of EtOAc once. The combined EtOAc-layers were dried (NaCl (saturated) and Na₂SO₄), filtered and concentrated. Separation by flash chromatography using *n*-heptane / EtOAc 4:1 yielded intermediate 42 and 0.24 g (65%) of intermediate 43 (E).

Example A7-1

Preparation of intermediate 33 and 40



Intermediate 33




Intermediate 40

2,6-dichloropurine (5.0 gram), methyl iodide (1.1 eq., 1.81 mL) and K₂CO₃ (1.2 eq., 4.39 gram) were dissolved in 200 mL of MeCN. After stirring for 70 hours at 20°C, the MeCN was evaporated. Both isomers were separated by column chromatography using CH₂Cl₂ / MeOH (90/10). Yield: 3.55 g (66%) of intermediate 33 and 1.61 g (30%) of intermediate 40.

-59-

Example A7-2

Preparation of intermediate 16 and 17




Intermediate 16

Intermediate 17

- 2,6-dichloro-1*H*-purine (18.32 g) and p-methoxybenzylchloride (1.0 eq., 15.72 g) were mixed with K₂CO₃ (1.0 eq, 15 g) in 100 ml of DMF. The reaction mixture was stirred vigorously and checked by TLC. The solvent was removed and the residue was
- 5 dissolved in 500 ml of EtOAc and 0.1 M NaOH (100 ml). The layers were separated. The EtOAc-layer was washed with 0.1 M NaOH (2x100 ml), dried (NaCl (saturated) and Na₂SO₄), filtered and concentrated. The residue was purified by column chromatography using n-heptane / EtOAc (1/1) as the eluent. The first fraction yielded 14.39 g of intermediate 16, the second fraction yielded 8.54 g of intermediate 17.

10 Example A8


a) Preparation of intermediate 18



- Intermediate 8 (prepared according to A4) (2.32 g) and intermediate 16 (prepared according to A7-2) (1.0 eq., 2.52 g) were mixed with K₂CO₃ (3.0 eq., 3.38 g) in 50 ml of DME. The reaction mixture was stirred vigorously at 80°C and checked by TLC and LC/MS. The organic solvents were removed and the residue was dissolved in 300 ml of
- 15 EtOAc and 100 ml of NaHCO₃ (saturated) (aqueous). The layers were separated. The water-layer was washed with 100 ml of EtOAc once, subsequently two times with 150 ml of CH₂Cl₂. The combined EtOAc-layers were dried (NaCl (saturated) and Na₂SO₄), filtered and concentrated. The combined CH₂Cl₂-layers were dried (NaCl (saturated) and Na₂SO₄), filtered and combined with the residue from the EtOAc-washings.
- 20 The residue was purified by trituration with 100 ml of ethanol. Yield : 2.69 g of intermediate 18 (72%).


-60-

b) Preparation of intermediate 20



Intermediate 19 (1.0 eq.; 5.81 g) (prepared according to A4) and intermediate 16 (prepared according to A7) (1.0 eq, 2.52 g.) were mixed with K_2CO_3 (3.0 eq; 3.38 g) in 100 ml of THF. The reaction mixture was stirred vigorously at 80°C and checked by TLC and LC/MS. The reaction mixture was dissolved in 400 ml of EtOAc and 100 ml of $NaHCO_3$ (saturated) (aqueous). The layers were separated. The water-layer was washed with 100 ml of EtOAc once. The combined EtOAc-layers were dried ($NaCl$ (saturated) and Na_2SO_4), filtered and concentrated. The residue was purified by trituration with ethanol. Yield: 0.24 g of intermediate 20 (54%).

c) Preparation of intermediate 22




Intermediate 21 (1.0 eq.; 290 mg) (prepared according to A4) and intermediate 16 (prepared according to A7) (1.0 eq, 2.52 g.) were mixed with K_2CO_3 (3.0 eq; 3.38 g) in 15 ml of t-amyl-OH. The reaction mixture was stirred vigorously at 80°C and checked by TLC and LC/MS. The organic solvents were removed and the residue was dissolved in 200 ml of EtOAc and 50 ml of $NaHCO_3$ (saturated) (aqueous). The layers were separated. The water-layer was washed with 50 ml of EtOAc once. The combined EtOAc-layers were dried ($NaCl$ (saturated) and Na_2SO_4), filtered and concentrated. The residue was purified by preparative TLC using n-heptane / $MeOCH_2CH_2OMe$ (2/3) as the eluent. Yield: 0.10 g of intermediate 22 (20%).

-61-


Example A9

a) Preparation of intermediate 23



Intermediate 16 (prepared according to A7) (50 mg) and intermediate 7 (prepared according to A3b) (5 eq., 144 mg) were dissolved in EtOH and water (3:1, 4 ml). The reaction mixture was stirred at 80°C and checked by TLC and LC/MS. The organic solvents were removed and the residue was dissolved in 100 ml of EtOAc and 50 ml of water. The layers were separated. The water-layer was washed with 50 ml of EtOAc once. The combined EtOAc-layers were washed with 1 M HCl (2x50 ml), 50 ml of NaHCO₃ (saturated) (aqueous), dried (NaCl (saturated) and Na₂SO₄), filtered and concentrated. The residue was purified by preparative TLC using n-heptane / EtOAc (1/1) as the eluent. Yield: 0.04 g of intermediate 23 (49%).

b) Preparation of intermediate 24




Intermediate 17 (prepared according to A7-2) (150 mg) and 4-ethoxy-2,6-dimethylbenzenamine (5 eq., 241 mg) were dissolved in EtOH and water (3:1, 8 ml). The reaction mixture was stirred at 80°C and checked by TLC and LC/MS. The organic solvents were removed and the residue was dissolved in 150 ml of EtOAc and 50 ml of water. The layers were separated. The water-layer was washed with 50 ml EtOAc once. The combined EtOAc-layers were washed with 1 M HCl (2x50 ml), 50 ml of NaHCO₃ (saturated) (aqueous), dried (NaCl (saturated) and Na₂SO₄), filtered and concentrated. The residue was purified by preparative TLC using n-heptane / MeOCH₂CH₂OMe (2/3) as the eluent. Yield: 0.15 g of intermediate 24 (63%).

-62-


Example A10

a) Preparation of intermediate 26




- Intermediate 25 (prepared according to A2) (113 mg) was suspended in 10 ml of THF at 20°C and stirred. KOtBu (1.0 eq., 60 mg) was added at once. After stirring for 30 minutes at 20°C, a solution of intermediate 17 (prepared according to A7-2) (1.0 eq., 150 mg) in 10 ml of THF was added dropwise. The reaction mixture was stirred at 20°C and checked by TLC and LC/MS. The organic solvents were removed and the residue was dissolved in 15 ml of CH₂Cl₂ and 5 ml of NaHCO₃ (saturated) (aqueous). The layers were separated. The CH₂Cl₂-layer was dried (Na₂SO₄), filtered and concentrated. The residue was purified by preparative TLC using n-heptane / MeOCH₂CH₂OMe (2/3) as the eluent. Yield: 0.14 g of intermediate 26 (46%).

b) Preparation of intermediate 28




-63-



Intermediate 27 (prepared according to A2) (84.5 mg) was suspended in 10 ml of THF at 20°C and stirred. KOtBu (1.0 eq., 36.3 mg) was added at once. After stirring for 30 minutes at 20°C, a solution of intermediate 16 (prepared according to A7-2) (1.0 eq., 100 mg) in 10 ml of THF was added dropwise. The reaction mixture was stirred at 20°C and checked by TLC and LC/MS. The reaction-mixture was dissolved in 100 ml of EtOAc and 50 ml of NaHCO₃ (saturated) (aqueous). The layers were separated. The water-layer was washed with 50 ml of EtOAc once. The combined EtOAc-layers were dried (NaCl (saturated) and Na₂SO₄), filtered and concentrated. The residue was triturated with diisopropyl ether, filtered and air-dried. Yield: 0.17 g of intermediate 28 (96%).

c) Preparation of intermediate 29




4-Hydroxy-3,5-dimethylbenzonitrile (143 mg) and intermediate 16 (prepared according to A7-2) (1.0 eq., 300 mg) were mixed with K₂CO₃ (3.0 eq, 402 mg) in 10 ml of 2-BuOH. The reaction mixture was stirred at 100°C and checked by TLC and LC/MS. The organic solvents were removed and the residue was dissolved in 100 ml of EtOAc and 50 ml of NaHCO₃ (saturated) (aqueous). The layers were separated. The water-layer was washed with 50 ml of EtOAc once. The combined EtOAc-layers were dried (NaCl (saturated) and Na₂SO₄), filtered and concentrated. The residue was purified by trituration with 10 ml of ethanol. Yield: 0.26 g of intermediate 29 (64%).

-64-

Example A11

Preparation of intermediate 31




Intermediate 30 (100 mg)(prepared according to A10a) was dissolved in 3 ml POCl₃ at 20°C. The reaction mixture was stirred at 20°C and checked by TLC. The reaction mixture was added dropwise to 200 ml of diisopropyl ether. The precipitate was filtered off and was dissolved in 200 ml of EtOAc and 200 ml of NaHCO₃ (saturated) (aqueous). The layers were separated. The water-layer was washed with 50 ml of EtOAc once. The combined EtOAc-layers were dried (NaCl (saturated) and Na₂SO₄), filtered and concentrated. Yield: 0.08 g of intermediate 31 (87%).

-65-

Example A12


Preparation of intermediate 32




Intermediate 16 (prepared according to A7-2) (3.09 gram), 4-cyanoaniline (1.3 eq., 1.54 gram) and collidine (2.3 eq., 3.03 ml) were dissolved in EtOH and water (3:1, 150 ml). The reaction mixture was stirred at 85°C and checked by TLC and LC/MS. After 150 hours, the reaction was cooled to ambient temperature and the solid was filtered off. Yield: 1.84 g of intermediate 32(47%).

Example A13

a. Preparation of intermediate 34 and 35



Intermediate 34




Intermediate 35

2-Amino-6-chloropurine (5.88 gram), methyl iodide (1.0 eq., 2.39 mL) and K_2CO_3 (1.0 eq., 4.88 gram) were dissolved in 100 ml of DMF. After stirring for 20 hours at 20°C, the solid material was filtered off and the DMF was evaporated. The residue was used for column chromatography using CH_2Cl_2 / MeOH (95/5). Yield: 2.77 gr (43%) of intermediate 34 and a mix-fraction. This mix-fraction was used for flash chromatography using CH_2Cl_2 /MeOH 95/5, slowly raised to 9/1. Yield: 0.39 gram (6%) of intermediate 34 and 0.60 g (9%) of intermediate 35. Total yield of intermediate 34: 3.16 g (49%), total yield of intermediate 35: 0.60 g (9%).

-66-


b. Preparation of intermediate 36



- Intermediate 34 (prepared according to A13.a) (370 mg), 4-iodobenzonitrile (1.3 eq, 600 mg), Pd(OAc)₂ (0.15 eq., 68 mg) and BINAP (0.3 eq., 380 mg) were mixed with Cs₂CO₃ (1.4 eq, 920 mg) in 35 ml of dioxane and N₂ was bubbled through the suspension for at least 20 minutes. The reaction mixture was stirred vigorously at 95°C in a sealed reaction-vessel for 48 hours and checked by TLC and LC/MS. Then the dioxane was evaporated and the residue was purified by column chromatography using CH₂Cl₂/MeOH 99/1. Yield: 151 mg of intermediate 36 (26%).


Example A14

a. Preparation of intermediate 37 and 38



- 2-Amino-6-chloropurine (5.47 gram), *p*-methoxy benzylchloride (1.0 eq., 5.05 gram) and K₂CO₃ (1.1 eq., 4.90 gram) were dissolved in 75 ml of DMF. After stirring for 20 hours at 20°C, the solid material was filtered off and the DMF was evaporated. Both isomers were separated by trituration with EtOAc. Yield: 6.73 gram (72%) of intermediate 37 and 0.48 gram (5%) of intermediate 38.

b. Preparation of intermediate 39



- Intermediate 37 (prepared according to A14.a) (1.86 gram), 4-bromobenzonitrile (1.3 eq, 1.52 gram), Pd(OAc)₂ (0.03 eq., 43 mg) and BINAP (0.06 eq., 240 mg) were mixed with Cs₂CO₃ (1.4 eq, 2.93 gram) in 370 ml of toluene and N₂ was bubbled through the suspension for at least 20 minutes. The reaction mixture was stirred vigorously at 80°C


-67-

in a sealed reaction-vessel for 80 hours and checked by TLC and LC/MS. Then the toluene was evaporated and the residue was purified by column chromatography using CH_2Cl_2 / MeOH (97.5/2.5). Yield: 0.67 gram of intermediate 39 (27%).

5 B. Preparation of the final compounds


Example B1

a) Preparation of compound 1



- Intermediate 26 (prepared according to A10a) (117 mg), 4-cyanoaniline (1.5 eq., 48 mg), $\text{Pd}(\text{OAc})_2$ (0.06 eq., 6.1mg) and BINAP (0.12 eq., 34 mg) were mixed with Cs_2CO_3 (1.2 eq, 107 mg) in 20 ml of toluene and N_2 was bubbled through the suspension for at least 20 minutes. The reaction mixture was stirred vigorously at 80°C in a sealed reaction-vessel and checked by TLC and LC/MS. The reaction mixture was dissolved in 150 ml of EtOAc and 50 ml of NaHCO_3 (saturated) (aqueous). The layers were separated. The water-layer was washed with 50 ml of EtOAc once. The combined EtOAc-layers were dried (NaCl (saturated) and Na_2SO_4), filtered and concentrated.
- 15 The residue was purified by preparative TLC using n-heptane / $\text{MeOCH}_2\text{CH}_2\text{OMe}$ (2/3) as the eluent. Yield : 0.11 g of compound 1 (82%).


b) Preparation of compound 2



-68-

Intermediate 18 (prepared according to A8a) (2.69 g), 4-cyanoaniline (1.5 eq., 1.03 g), Pd(OAc)₂ (0.05 eq., 66 mg) and BINAP (0.10 eq., 363 mg) were mixed with Cs₂CO₃ (1.2 eq, 2.47 g) in 75 ml of toluene and argon was bubbled through the suspension for at least 20 minutes. The reaction mixture was stirred vigorously at 80°C in a sealed reaction-vessel and checked by TLC and LC/MS. The reaction mixture was filtered off. The residue was washed once with 25 ml of toluene and dissolved in 300 ml of EtOAc and 100 ml of NaHCO₃ (saturated) (aqueous). The layers were separated. The EtOAc-layer was dried (Na₂SO₄), filtered and concentrated. The residue was dissolved in 50 ml of CH₂Cl₂/MeOH (25:1), followed by addition of 500 ml of diisopropylether. The precipitate was filtered off and air-dried. Yield : 2.05 g of compound 2 (65%).


c) Preparation of compound 3



Intermediate 20 (prepared according to A8b) (242 mg), 4-cyanoaniline (1.5 eq.; 96 mg), Pd₂(dba)₃ (0.03 eq.; 15 mg) and BINAP (0.06 eq.; 20 mg) were mixed with Cs₂CO₃ (1.2 eq; 212 mg) in 20 ml of toluene and N₂ was bubbled through the suspension for at least 20 minutes. The reaction mixture was stirred at 80°C in a sealed reaction-vessel and checked by TLC and LC/MS. The reaction mixture was dissolved in 150 ml of EtOAc and 50 ml of NaHCO₃ (saturated) (aqueous). The layers were separated. The water-layer was washed with 50 ml of EtOAc once. The combined EtOAc-layers were dried (NaCl (saturated) and Na₂SO₄), filtered and concentrated. The residue was purified by preparative TLC using n-heptane / MeOCH₂CH₂OMe (2/3) as the eluent. Yield : 0.13 g of compound 3 (46%).


-69-

d) Preparation of compound 4



Intermediate 31 (prepared according to A11) (80 mg), 4-cyanoaniline (1.5 eq.; 32 mg), $\text{Pd}_2(\text{dba})_3$ (0.07 eq.; 12 mg) and BINAP (0.14 eq.; 16 mg) were mixed with Cs_2CO_3 (1.2 eq.) in 20 ml of toluene and N_2 was bubbled through the suspension for at least 20 minutes. The reaction mixture was stirred at 80°C in a sealed reaction-vessel and checked by TLC and LC/MS. The reaction mixture was dissolved in 150 ml of EtOAc and 50 ml of NaHCO_3 (saturated) (aqueous). The layers were separated. The water-layer was washed with 50 ml of EtOAc once. The combined EtOAc-layers were dried (NaCl (saturated) and Na_2SO_4), filtered and concentrated. The residue was purified by preparative TLC using n-heptane / $\text{MeOCH}_2\text{CH}_2\text{OMe}$ (2/3) as the eluent. Yield : 0.08 g of compound 4 (84%).

e) Preparation of compound 73




Intermediate 32 (prepared according to A12) (150 mg), intermediate 6 (prepared according to A3.a) (1.5 eq., 99 mg), $\text{Pd}(\text{OAc})_2$ (0.07 eq., 6 mg) and BINAP (0.14 eq., 33 mg) were mixed with Cs_2CO_3 (1.2 eq, 150 mg) in 15 ml of toluene and N_2 was bubbled through the suspension for at least 20 minutes. The reaction mixture was stirred vigorously at 80°C in a sealed reaction-vessel and checked by TLC and LC/MS. The reaction mixture was dissolved in 150 ml of EtOAc and 50 ml of NaHCO_3 (saturated) (aqueous). The layers were separated. The water-layer was washed with 50 ml of EtOAc once. The combined EtOAc-layers were dried (NaCl (saturated) and

-70-

Na₂SO₄), filtered and concentrated. The residue was purified by preparative TLC using CH₂Cl₂/MeOH (95/5) as the eluent. Yield : 0.12 g of compound 73 (59%).

f) Preparation of compound 57




Intermediate 36 (40 mg) (prepared according to A13.b), intermediate 43 (1.3 eq., 34 mg) (prepared according to A6-1.b), Pd(OAc)₂ (0.20 eq., 6.3 mg) and BINAP (0.40 eq., 35 mg) were mixed with Cs₂CO₃ (1.3 eq, 60 mg) in 25 ml of dioxane and argon was bubbled through the suspension for at least 40 minutes. The reaction mixture was stirred vigorously at 95°C in a sealed reaction-vessel. After 20 hours, the reaction mixture was concentrated. The residue was used for flash chromatography using CH₂Cl₂/MeOH 99:1. Yield : 23 mg of compound 57 (38%).

10

Example B2

a) Preparation of compound 5




Compound 1 (100 mg) was dissolved in 2 ml of TFA. The reaction mixture was stirred at 40 °C and checked by TLC and LC/MS. The reaction mixture was diluted with 150 ml of CH₂Cl₂ and added dropwise to a concentrated aqueous solution of K₂CO₃ (200 ml), checking pH continuously. Extra CH₂Cl₂ (100 ml) was added to dissolve all material. The layers were separated. The CH₂Cl₂-layer was dried (NaCl (saturated) and

15

-71-


Na₂SO₄), filtered and concentrated till around 5 ml. The precipitated solid was filtered off and washed once with 5 ml of CH₂Cl₂. Yield : 0.01 g of compound 5 (15%).

b) Preparation of compound 6



Compound 2 (1.79 g) was dissolved in 8 ml of TFA. The reaction mixture was stirred at 60°C and checked by TLC and LC/MS. The reaction mixture was added dropwise to a 2M aqueous solution of NaOH (400 ml), checking pH continuously. EtOAc (800 ml) was added to dissolve all material. The layers were separated. The EtOAc-layer was dried (NaCl (saturated) and Na₂SO₄), filtered and concentrated. The residue was purified by column chromatography using CH₂Cl₂/ MeOH (95/5) as the eluent. The product was isolated by trituration with 500 ml of diisopropylether. The precipitate was filtered off and air-dried. Yield: 0.71 g of compound 6 (51%).

c) Preparation of compound 70




Compound 73 (prepared according to B1.e) (108 mg) was dissolved in 2 ml of TFA. The reaction mixture was stirred at 60°C and checked by TLC and LC/MS. The reaction mixture was diluted with 150 ml of CH₂Cl₂ and added dropwise to a concentrated aqueous solution of K₂CO₃ (200 ml), checking pH continuously. Extra CH₂Cl₂ (100 ml) was added to dissolve all material. The layers were separated. The CH₂Cl₂-layer was dried (NaCl (saturated) and Na₂SO₄), filtered and concentrated till around 5 ml and purified by preparative TLC using CH₂Cl₂/MeOH (95/5) as the eluent. Yield : 0.034 g of compound 70 (44%).

-72-


Example B3

a) Preparation of compound 49



Compound 6 (prepared according to B2.b) (30 mg) was dissolved in 20 ml of MeCN, and isopropyl iodide (1.0 eq., 12 mg) and K_2CO_3 (3 eq, 29 mg) were added. Stirring for 17 days at 20°C. The MeCN was evaporated and the residue was purified by preparative TLC using $CH_2Cl_2/MeOH$ (9/1) as the eluent. Yield : 12 mg of compound 49 (34%).


b) Preparation of compound 60



Compound 6 (prepared according to B2.b) (30 mg) was dissolved in 20 ml of MeCN, and iodo acetonitrile (1.0 eq.) and K_2CO_3 (3 eq, 29 mg) were added. Stirring for 17 days at 20°C. The MeCN was evaporated and the residue was purified by preparative TLC using $CH_2Cl_2/MeOH$ (9/1) as the eluent. Yield : compound 60 (52 %).

-73-

c) Preparation of compound 48




Compound 6 (prepared according to B2.b) (30 mg) was dissolved in 20 ml of MeCN, and ethyl iodide (1.0 eq.) and K_2CO_3 (3 eq, 29 mg) were added. Stirring for 17 days at 20°C. The MeCN was evaporated and the residue was purified by preparative TLC using $CH_2Cl_2/MeOH$ (9/1) as the eluent. Yield : compound 48 (67 %).

5

Example B4

a) Preparation of compound 75




Final compound 6 (prepared according to B2.b)(214 mg) was dissolved in 60 ml of MeCN and 2-bromo methyl acetate (1.0 eq., 48 microliter) and K_2CO_3 (3 eq, 210 mg) were added. Stirring for 100 hours at 20°C. The MeCN was evaporated and the residue was purified by Flash chromatography using $CH_2Cl_2/MeOH$: 99/1 to 96/4 as the gradient eluent. Yield :139 mg of compound 75 (56%).

10

-74-


b. Preparation of compound 50



Compound 75 (prepared according to B4a) (50 mg) was dissolved in 10 ml of THF and 50 ml of EtOH and 3 portions of 0.20 gram of NaBH₄ were added initially and every half hour. Stirring for 2 hours at 0°C. The reaction mixture was diluted with EtOAc and washed with 1 N HCl, with saturated aqueous NaHCO₃ and dried (NaCl (saturated) and Na₂SO₄), filtered and concentrated and the residue was purified by preparative TLC using CH₂Cl₂/MeOH (95/5) as the eluent.. Yield :15 mg of compound 50 (32%).

Example B5


a. Preparation of compound 51



Compound 75 (prepared according to B4a) (15 mg) was dissolved in 2 ml of THF and NH₃ 7 N in MeOH (1 ml) was added. Stirring for 100 hours at 20°C. The solvents were evaporated and the residue was stirred in *i*-Pr₂O and filtered off. Yield :15 mg of compound 51 (99%).

-75-


b. Preparation of compound 52



Compound 51 (prepared according to B5.a) (27 mg) was dissolved in 2 ml of THF and POCl₃ (4 ml) was added. Stirring for 120 hours at 50°C. The reaction mixture was diluted with *i*-Pr₂O; an oil was formed on the bottom of the flask. The *i*-Pr₂ was decanted and the oil was stirred with 2 N NaOH and EtOAc. The EtOAc fraction was dried (NaCl (saturated) and Na₂SO₄), filtered and concentrated and the residue was purified by preparative TLC using CH₂Cl₂/MeOH (95/5) as the eluent.. Yield :15 mg of compound 52 (60%).

Tables 1 to 4 list the compounds of formula (I) which were prepared according to one of the above examples.

Table 1:



Co. no.	Exp. no.	R ²	R ³	R ^{4a}	R ^{4b}	X ₁	Physico-chem data.
7	B1c	CN	CH ₃	CH ₃	CH ₃	O	
8	B1b	CN	CH ₃	CH ₃	CH ₃	NH	
9	B1c	CN	C(CH ₃) ₃	CH ₃	CH ₃	NH	
10	B1c	CN	OCH ₃	OCH ₃	OCH ₃	NH	

-76-













Co. no.	Exp. no.	R ²	R ³	R ^{4a}	R ^{4b}	X ₁	Physico- chem data.
11	B1c	CN	CN	CH ₃	CH ₃	O	
12	B1c	CN	CN	CH ₃	CH ₃	NH	
13	B1c	CN		CH ₃	CH ₃	NH	
3	B1c	CN		CH ₃	CH ₃	NH	
14	B1b	CN		Cl	CH ₃	NH	
2	B1b	CN		OCH ₃	CH ₃	NH	
40	B1.f	CN		CH ₃	CH ₃	NH	
41	B1.f	CN		CH ₃	CH ₃	NH	(Z)
42	B1.f	CN		CH ₃	CH ₃	NH	(E)
43	B1.f	CN		OCH ₃	CH ₃	NH	
15	B1b	CN		CH ₃	CH ₃	NH	
16	B1c	CN		CH ₃	CH ₃	O	
17	B1c	CN		CH ₃	CH ₃	NH	


Table 2 :



-77-





















Co no.	Exp. no.	R ³	R ^{4a}	R ^{4b}	R ¹⁷	X ₁	Physico chem data.
18	B1a		CH ₃	CH ₃		NH	
44	B1.b		CH ₃	CH ₃	CH ₃	NH	
4	B1d		CH ₃	CH ₃		O	
19	B1d		OCH ₃	OCH ₃		O	
20	B1d		CH ₃	CH ₃		O	
21	B1d		CH ₃	CH ₃		O	
1	B1a		CH ₃	CH ₃		O	(E)

Table 3:



Co. no.	Exp. no.	R ²	R ³	R ^{4a}	R ^{4b}	R ¹⁷	X ₁	Physico chem data
22	B2b	CH ₃	CH ₃	CH ₃	CH ₃	H	NH	
23	B2b	CN	CH ₃	CH ₃	CH ₃	H	O	

-78-

Co. no.	Exp. no.	R ²	R ³	R ^{4a}	R ^{4b}	R ¹⁷	X ₁	Physico chem data
24	B2b	CN	CH ₃	CH ₃	CH ₃	H	NH	
45	B1f	CN	CH ₃	CH ₃	CH ₃	CH ₃	NH	
25	B2b	CN	OCH ₃	CH ₃	CH ₃	H	NH	
26	B2b	CN	OCH ₃	OCH ₃	OCH ₃	H	NH	
27	B2b	CN	C(CH ₃) ₃	CH ₃	CH ₃	H	NH	
28	B2b	CN	CN	CH ₃	CH ₃	H	O	
29	B2b	CN	CN	CH ₃	CH ₃	H	NH	
46	B1.b	CN	CN	CH ₃	CH ₃	CH ₃	NH	
30	B2a	CN		CH ₃	CH ₃	H	NH	
31	B2b	CN		CH ₃	CH ₃	H	NH	
32	B2b	CN		CH ₃	CH ₃	H	NH	
33	B2a	CN		CH ₃	CH ₃	H	O	
34	B2b	CN		CH ₃	CH ₃	H	NH	
47	B1.b	CN		CH ₃	CH ₃	CH ₃	NH	
6	B2b	CN		OCH ₃	CH ₃	H	NH	
48	B3c	CN		OCH ₃	CH ₃		NH	
49	B3.a	CN		OCH ₃	CH ₃		NH	
50	B4.b	CN		OCH ₃	CH ₃		NH	
51	B5.a	CN		OCH ₃	CH ₃		NH	
52	B5.b	CN		OCH ₃	CH ₃		NH	
35	B2a	CN		OCH ₃	OCH ₃	H	O	
53	B2.b	CN		OCH ₃	Cl	H	NH	
54	B2.b	CN		Cl	CH ₃	H	NH	(E)

-79-

Co. no.	Exp. no.	R ²	R ³	R ^{4a}	R ^{4b}	R ¹⁷	X ₁	Physico chem data
55	B2.b	CN		Cl	Cl	H	NH	(E)
56	B2.b	CN		CH ₃	CH ₃	H	NH	
57	B1.f	CN		CH ₃	CH ₃	CH ₃	NH	(E)
58	B2.b	CN		CH ₃	CH ₃	H	NH	(E)
59	B1.f	CN		CH ₃	CH ₃	CH ₃	NH	
60	B3.b	CN		CH ₃	CH ₃		NH	(E)
61	B2.b	CN		CH ₃	CH ₃	H	NH	(Z)
62	B2.b	CN		CH ₃	CH ₃	H	NH	
63	B2.b	CN		OCH ₃	CH ₃	H	NH	
64	B1.b	CN		CH ₃	CH ₃	CH ₃	NH	
65	B2.b	CN		CH ₃	CH ₃	H	NH	(E)
66	B1.f	CN		CH ₃	CH ₃	CH ₃	NH	(E)
36	B2.a	CN		CH ₃	CH ₃	H	O	
37	B2.b	CN		CH ₃	CH ₃	H	NH	
67	B2.a	CN		CH ₃	CH ₃	H	O	
5	B2.a	CN		CH ₃	CH ₃	H	O	

-80-












Co. no.	Exp. no.	R ²	R ³	R ^{4a}	R ^{4b}	R ¹⁷	X ₁	Physico chem data
38	B2b	CN		CH ₃	CH ₃	H	NH	
39	B2b	CN		CH ₃	CH ₃	H	O	
68	B1.b			CH ₃	CH ₃	CH ₃	NH	(E)
75	B4.a	CN		OCH ₃	CH ₃	CH ₂ -C(=O)-OCH ₃	NH	

Table 4:



Co. no.	Exp. no.	R ^{2a}	R ¹⁷	Physico chem data
69	B2.c	CN	H	
70	B2.c		H	
71	B1.e	CH ₃		
72	B1.e	CN		
73	B1.e			
74	B2c	CH ₃	H	

C. Analytical data

Table 5 lists ¹H NMR data for the compounds of formula (I). All spectra were recorded (300 MHz) in DMSO-d₆, unless otherwise specified. The shifts (δ) are in ppm, relative to TMS. Between parenthesis, the number of H's, the peak-form and the coupling (J) in Hz are indicated. The following abbreviations were used : s :singlet, bs : broad singlet, ds : double singlet, d : doublet, dd : double doublet, t : triplet, dt : double triplet, q : quartet, dq : double quartet, h : heptet, m : multiplet.

Table 5

10

Co.No.	NMR results
67	δ = 13.23 (1, bs), 8.23 (1, s), 7.61 (2, d, 9), 7.60 (2, s), 7.50 (1, d, 15), 7.40 (2, d, 9), 7.23 (1, d, 15), 2.95 & 3.20 (6, ds), 2.11 (6, s)
54	δ = 12.70 (1, bs), 9.46 (2, bs), 7.98 (1, s), 7.79 (1, bs), 7.50-7.75 (4, m), 7.39 (2, d, 9), 6.59 (1, d, 17), 2.24 (3, s)
55	δ = 12.74 (1, bs), 9.77(1, bs), 9.51(1, bs), 7.99 (1, bs), 7.95 (2, s), 7.60-7.80 (3, m), 7.42 (2, d, 9), 6.72 (1, d, 17)
45	δ = 9.53 (1, bs), 9.13 (1, bs), 7.95 (1, s), 7.66 (2, d, 8), 7.31 (2, d, 8), 6.97 (2, s), 3.69 (3, s), 2.33 (3, s), 2.10 (6, s)
44	δ = 9.41(1, bs), 8.44(1, bs), 8.12 (1, bs), 7.69 (1, d, 17), 7.50-7.60 (4, m), 7.28 (2, d, 9), 6.51 (1, d, 17), 4.10 (3, s), 2.21 (6, s)
70	δ = 12.55 (1, bs), 9.99 (1, bs), 8.44 (1, bs), 8.00 (2, bs), 7.90 (1, bs), 7.62 (1, d, 17), 7.44 (2, s), 7.44 (2, bs), 6.41 (1, d, 17), 2.18 (6, s)
73	δ = 10.06 (1, bs), 8.60 (1, bs), 8.03 (1, s), 7.89 (2, bs), 7.64 (1, d, 17), 7.47 (2, s), 7.40 (2, bs), 7.23 (2, d, 9), 6.89 (2, d, 9), 6.44 (1, d, 17), 5.15 (2, bs), 3.73 (3, s), 2.16 (6, s)
47	δ = 9.58 (1, bs), 9.37 (1, bs), 7.97 (1, s), 7.60-7.70 (3, m), 7.48 (2, s), 7.33 (2, d, 9), 6.47 (1, d, 17), 3.70 (3, s), 2.17 (6, s)
53	δ = 12.65 (1, bs), 9.46 (1, bs), 9.24 (1, bs), 7.96 (1, bs), 7.60-7.80 (3, m), 7.52 (1, s), 7.30-7.48 (3, m), 6.69 (1, d, 17), 3.77 (3,s)
46	δ = 9.61 (1, bs), 9.51 (1, bs), 7.99 (1, s), 7.69 (2, s), 7.64 (2, d, 8), 7.38 (2, d, 8), 3.70 (3, s), 2.20 (6, s)

-82-

Co.No.	NMR results
63	$\delta = 12.61$ (1, bs), 9.41 (1, bs), 8.93 (1, bs), 7.94 (1, s), 7.70 (2, d, 8), 7.30-7.50 (3, m), 7.05 (2, s), 3.70 (3, s), 2.20 (6, s)
43	$\delta = 9.49$ (1, bs), 9.04 (1, bs), 8.08 (1, s), 7.70 (2, d, 8), 7.35-7.50 (3, m), 7.31 (2, d, 9), 7.05 (1, s), 7.04 (1, s), 6.92 (2, d, 9), 5.25 (2, s), 3.71 (3, s), 3.70 (3, s), 2.20 (6, s)
62	$\delta = 12.64$ (1, bs), 9.41 (1, bs), 9.25 (1, bs), 7.97 (1, s), 7.69 (2, d, 8), 7.30-7.45 (3, m), 7.28 (2, s), 2.19 (9, m)
42	$\delta = 9.49$ (1, bs), 9.35 (1, bs), 8.10 (1, s), 7.69 (2, d, 8), 7.30-7.45 (5, m), 7.27 (2, s), 6.92 (2, d, 9), 5.25 (2, s), 3.72 (3, s), 2.18 (9, s)
71	$\delta = 10.00$ (1, bs), 7.80-8.50 (4, m), 7.20-7.40 (4, m), 6.95 (2, s), 6.88 (2, d, 8), 5.15 (2, bs), 3.73 (3, s), 2.32 (3, s), 2.10 (6, s)
61	$\delta = 12.66$ (1, bs), 9.43 (1, bs), 9.27 (1, bs), 7.96 (1, s), 7.70 (2, d, 8), 7.50 (2, s), 7.39 (2, d, 8), 7.27 (1, s), 2.19 (6, s), 2.17 (3, s)
41	$\delta = 9.50$ (1, bs), 9.36 (1, bs), 8.10 (1, s), 7.70 (2, d, 8), 7.49 (2, s), 7.40 (2, d, 8), 7.33 (2, d, 9), 7.27 (1, s), 6.92 (2, d, 9), 5.25 (2, s), 3.72 (3, s), 2.18 (6, s), 2.16 (3, s)
69	$\delta = 12.60$ (1, bs), 10.04 (1, bs), 8.59 (1, bs), 8.01 (2, d, 8), 7.93 (1, s), 7.62 (2, s), 7.52 (2, d, 8), 2.20 (6, s)
72	$\delta = 10.11$ (1, bs), 8.75 (1, bs), 8.05 (1, s), 7.89 (2, d, 8), 7.66 (2, s), 7.45 (2, d, 8), 7.22 (2, d, 9), 6.89 (2, d, 9), 5.15 (2, s), 3.73 (3, s), 2.18 (6, s)
64	$\delta = 9.58$ (1, bs), 9.34 (1, bs), 7.97 (1, s), 7.65 (2, d, 8), 7.43 (1, s), 7.33 (2, d, 8), 7.29 (2, s), 3.70 (3, s), 2.18 (9, s)
51	$\delta = 9.53$ (1, bs), 9.08 (1, bs), 7.92 (1, s), 7.60-7.70 (4, m), 7.00-7.40 (5, m), 6.57 (1, d, 17), 4.75 (2, s), 3.73 (3, s), 2.20 (3, s)
52	$\delta = 9.66$ (1, bs), 9.26 (1, bs), 8.05 (1, s), 7.60-7.75 (3, m), 7.00-7.40 (4, m), 6.58 (1, d, 17), 5.36 (2, s), 3.72 (3, s), 2.19 (3, s)
50	$\delta = 9.53$ (1, bs), 9.06 (1, bs), 7.94 (1, s), 7.60-7.75 (3, m), 7.25-7.40 (3, m), 7.22 (1, s), 6.57 (1, d, 16), 5.01 (1, t, 5), 4.14 (2, t, 5), 3.70-3.80 (5, m), 2.19 (3, s)

-83-

Co.No.	NMR results
48	$\delta = 9.54$ (1, bs), 9.06 (1, bs), 8.02 (1, s), 7.60-7.75 (3, m), 7.25-7.40 (3, m), 7.21 (1, s), 6.57 (1, d, 17), 4.13 (2, q, 7), 3.72 (3, s), 2.18 (3, s), 1.43 (3, t, 7)
49	$\delta = 9.49$ (1, bs), 9.05 (1, bs), 8.10 (1, s), 7.60-7.75 (3, m), 7.25-7.40 (3, m), 7.21 (1, s), 6.56 (1, d, 17), 4.67 (1, h, 7), 3.72 (3, s), 2.19 (3, s), 1.55 (6, d, 7)
56	$\delta = 12.66$ (1, bs), 9.41 (1, bs), 9.26 (1, bs), 7.96 (1, bs), 7.71 (2, d, 8), 7.46 (2, s), 7.38 (2, d, 8), 6.14 (1, s), 2.46 (3, s), 2.20 (6, s)
40	$\delta = 9.48$ (1, bs), 9.35 (1, bs), 8.11 (1, bs), 7.71 (2, d, 8), 7.45 (2, s), 7.39 (2, d, 8), 7.34 (2, d, 9), 6.92 (2, d, 9), 6.14 (1, s), 5.25 (2, s), 3.71 (3, s), 2.46 (3, s), 2.18 (6, s)
58	$\delta = 12.65$ (1, bs), 9.42 (1, bs), 9.26 (1, bs), 7.96 (1, bs), 7.67 (2, d, 8), 7.42 (2, s), 7.35 (2, d, 8), 6.03 (1, s), 2.90 (2, q, 8), 2.20 (6, s), 1.12 (3, t, 8)
57	$\delta = 9.56$ (1, bs), 9.34 (1, bs), 7.97 (1, bs), 7.67 (2, d, 8), 7.47 (2, s), 7.33 (2, d, 8), 6.15 (1, s), 3.70 (3, s), 2.47 (3, s), 2.19 (6, s)
59	$\delta = 9.59$ (1, bs), 9.34 (1, bs), 7.97 (1, bs), 7.64 (2, d, 8), 7.43 (2, s), 7.31 (2, d, 8), 6.03 (1, s), 3.70 (3, s), 2.91 (2, q, 8), 2.19 (6, s), 1.12 (3, t, 8)
65	$\delta = 12.64$ (1, bs), 9.43 (1, bs), 9.20 (1, bs), 7.96 (1, bs), 7.67 (2, d, 8), 7.36 (2, d, 8), 7.10 (2, s), 2.37 (3, s), 2.19 (6, s), 1.89 (3, s)
66	$\delta = 9.59$ (1, bs), 9.28 (1, bs), 7.97 (1, bs), 7.64 (2, d, 8), 7.32 (2, d, 8), 7.11 (2, s), 3.70 (3, s), 2.38 (3, s), 2.18 (6, s), 1.88 (3, s)
68	$\delta = 9.22$ (2, bs), 7.92 (1, s), 7.45-7.70 (9, m), 7.01 (1, d, 16), 3.69 (3, s), 2.18 (6, s)
60	$\delta = 9.67$ (1, bs), 9.52 (1, bs), 8.08 (1, bs), 7.67 (2, d, 8), 7.44 (2, s), 7.33 (2, d, 8), 6.04 (1, s), 5.37 (2, s), 2.91 (2, q, 8), 2.20 (6, s), 1.12 (3, t, 8)

-84-

Co.No.	NMR results
1	δ = 9.81 (H, bs), 8.66 (H, s), 7.57 (2H, d, 8), 7.54 (2H, s), 7.43-7.50 (H, m), 7.36 (2H, d, 8), 7.20 (2H, d, 8), 7.16 (H, d, 16), 6.90 (2H, d, 8), 5.57 (2H, s), 3.71 (3H, s), 3.57 & 3.45 (2H, dq, 7), 3.15 & 2.93 (3H, ds), 1.89 (6H, s), 1.17 & 1.07 (3H, dt, 7)
2	δ = 9.50 (H, bs), 9.07 (H, bs), 8.08 (H, bs), 7.70 (2H, d, 9), 7.65 (H, d, 17), 7.40 (2H, d, 9), 7.31 (2H, d, 8), 7.30 (H, s), 7.20 (H, s), 6.91 (2H, d, 8), 6.55 (H, d, 17), 5.25 (2H, s), 3.71 (6H, s), 2.18 (3H, s)
3	δ = 9.52 (H, bs), 9.40 (H, bs), 8.12 (H, bs), 7.60-7.80 3H, m), 7.48 (2H, s), 7.30-7.43 (4H, m), 6.93 (2H, d, 8), 6.47 (H, d, 17), 5.26 (2H, s), 3.71 (3H, s), 2.17 (6H, s)
4	(CDCl ₃) δ = 8.08 (H, s), 7.51 (2H, d, 9), 7.39 (H, d, 17), 7.35 (2H, d, 9), 7.24 (2H, s), 7.20 (H, bs), 7.14 (2H, d, 9), 6.87 (2H, d, 9), 5.89 (H, d, 17), 5.51 (2H, s), 3.79 (3H, s), 1.97 (6H, s)
5	δ = 13.15 (H, bs), 9.86 (H, s), 8.23 (H, s), 7.59 (2H, s), 7.58 (2H, d, 8), 7.41-7.53 H, m), 7.37 (2H, d, 8), 7.18 (H, d, 15), 3.59 & 3.42 (2H, dq, 7), 3.16 & 2.93 (3H, ds), 2.11 (6H, s), 1.18 & 1.08 (3H, dt, 7)
6	δ = 12.61 (H, bs), 9.42 (H, bs), 8.95 (H, bs), 7.95 (H, bs), 7.71 (2H, d, 9), 7.65 (H, d, 17), 7.39 (2H, d, 9), 7.30 (H, s), 7.21 (H, s), 6.55 (H, d, 17), 3.73 (3H, s), 2.19 (3H, s)
7	(CDCl ₃) δ = 7.79 (1, s), 7.35-7.50 (4, m), 7.31 (2, d, 9), 7.17 (1, bs), 6.97 (2, s), 6.91 (2, d, 9), 5.27 (2, s), 3.81 (3, s), 2.38 (3, s), 2.13 (6, s).
8	δ = 9.47 (1, bs), 9.17 (1, bs), 8.10 (1, bs), 7.72 (2, d, 8), 7.30-7.45 (4, m), 6.90-7.00 (4, m), 5.25 (2, s), 3.72 (3, s), 2.31 (3, s), 2.10 (6, s)
9	(CDCl ₃) δ = 7.59 (1, bs), 7.52 (2, d, 8), 7.35 (2, d, 8), 7.27 (2, d, 9), 7.16 (2, s), 7.03 (1, bs), 6.93 (1, bs), 6.88 (2, d, 9), 5.21 (2, s), 3.79 (3, s), 2.25 (6, s), 1.38 (9, s)
10	(CDCl ₃) δ = 7.55-7.65 (3, m), 7.42 (2, d, 8), 7.26 (2, d, 9), 7.08 (1, bs), 6.89 (2, d, 9), 6.81 (1, bs), 6.25 (2, s), 5.21 (2, s), 3.89 (3, s), 3.80 (3, s), 3.76 (6, s)

-85-

Co.No.	NMR results
11	(CDCl ₃) δ = 7.83 (1, s), 7.45-7.55 (6, m), 7.32 (2, d, 9), 7.10 (1, bs), 6.92 (2, d, 9), 5.30 (2, s), 3.81 (3, s), 2.20 (6, s)
12	(CDCl ₃) δ = 7.50-7.65 (3, m), 7.40-7.50 (4, m), 7.20-7.35 (3, m), 7.06 (1, bs), 6.94 (2, d, 9), 5.25 (2, s), 3.81 (3, s), 2.29 (6, s)
13	(CDCl ₃) δ = 8.19 (1, bs), 7.82 (1, s), 7.54 (2, d, 8), 7.41 (2, d, 8), 7.20-7.30 (3, m), 7.04 (2, s), 6.88 (2, d, 9), 5.21 (2, s), 3.79 (3, s), 2.97 (2, t, 6), 2.69 (2, t, 6), 2.16 (6, s)
14	δ = 9.59 (1, bs), 9.54 (1, bs), 8.12 (1, bs), 7.60-7.80 (5, m), 7.30-7.45 (4, m), 6.92 (2, d, 9), 6.59 (1, d, 17), 5.26 (2, s), 3.71 (3, s), 2.23 (3, s)
15	(CDCl ₃) δ = 7.68 (1, d, 15), 7.64 (1, s), 7.56 (2, d, 8), 7.39 (2, d, 8), 7.35 (2, s), 7.31 (2, d, 9), 7.08 (2, bs), 6.85-6.95 (3, m), 5.24 (2, s), 3.81 (3, s), 3.22 (3, s), 3.10 (3, s), 2.28 (6, s)
16	(CDCl ₃) δ = 7.81 (1, s), 7.74 (1, d, 17), 7.40-7.50 (4, m), 7.27-7.35 (4, m), 7.16 (1, bs), 6.92 (2, d, 9), 6.85 (1, d, 17), 5.28 (2, s), 3.81 (3, s), 3.76 (8, bs), 2.18 (6, s)
17	δ = 9.58 (1, bs), 7.95 (2, d, 8), 7.40-7.65 (6, m), 7.29 (1, d, 17), 7.12 (2, d, 9), 6.96 (2, d, 9), 5.28 (1, s), 5.02 (2, s), 3.79 (3, s), 3.50-3.78 (8, m), 2.31 (6, s)
18	δ = 9.44 (1, bs), 8.36 (1, s), 7.88 (1, bs), 7.53 (2, d, 8), 7.28 (2, d, 8), 7.11 (2, d, 9), 6.92 (2, d, 9), 6.74 (2, s), 5.70 (2, s), 4.06 (2, q, 7), 3.71 (3, s), 1.89 (6, s), 1.36 (3, t, 7)
19	(CDCl ₃) δ = 7.99 (1, bs), 7.54 (2, d, 8), 7.45 (1, d, 17), 7.30-7.40 (4, m), 7.13 (1, bs), 6.87 (2, d, 9), 6.78 (2, s), 5.94 (1, d, 17), 5.50 (2, s), 3.80 (3, s), 3.77 (6, s)
20	δ = 9.83 (1, bs), 8.68 (1, s), 7.61 (2, d, 8), 7.51 (1, bs), 7.35-7.50 (5, m), 7.21 (2, d, 9), 7.11 (1, bs), 6.91 (2, d, 9), 6.62 (1, d, 17), 5.58 (2, s), 3.71 (3, s), 1.89 (6, s)
21	(CDCl ₃) δ = 8.07 (1, s), 7.68 (1, d, 17), 7.46 (2, d, 8), 7.30-7.36 (4, m), 7.24 (1, bs), 7.16 (2, d, 9), 6.85-6.95 (3, m), 5.52 (2, s), 3.80 (3, s), 3.23 (3, s), 3.10 (3, s), 1.98 (6, s)

-86-

Co.No.	NMR results
23	$\delta = 9.85$ (1, bs), 8.20 (1, bs), 7.62 (2, d, 8), 7.42 (2, d, 8), 7.03 (2, s), 2.34 (3, s), 2.04 (6, s)
24	$\delta = 12.64$ (1, bs), 9.39 (1, bs), 9.07 (1, bs), 7.96 (1, bs), 7.72 (2, d, 8), 7.38 (2, d, 8), 6.98 (2, s), 2.31 (3, s), 2.12 (6, s)
25	$\delta = 12.65$ (1, bs), 9.43 (1, bs), 9.04 (1, bs), 8.03 (1, s), 7.73 (2, d, 8), 7.39 (2, d, 8), 6.77 (2, s), 3.79 (3, s), 2.14 (6, s)
26	$\delta = 12.50$ (1, bs), 9.39 (1, bs), 8.37 (1, bs), 7.93 (1, bs), 7.76 (2, d, 8), 7.43 (2, d, 8), 6.38 (2, s), 3.86 (3, s), 3.69 (6, s)
27	$\delta = 12.64$ (1, bs), 9.42 (1, bs), 9.09 (1, bs), 7.97 (1, bs), 7.69 (2, d, 8), 7.35 (2, d, 8), 7.18 (2, s), 2.15 (6, s), 1.34 (9, s)
28	$\delta = 13.24$ (1, bs), 9.90 (1, bs), 8.27 (1, s), 7.80 (2, s), 7.60 (2, d, 8), 7.48 (2, d, 8), 2.14 (6, s)
29	$\delta = 12.73$ (1, bs), 9.47 (2, bs), 8.01 (1, s), 7.65 - 7.75 (4, m), 7.44 (2, d, 8), 2.21 (6, s)
30	$\delta = 12.73$ (1, bs), 9.41 (1, bs), 9.02 (1, bs), 8.03 (1, bs), 7.69 (2, d, 8), 7.36 (2, d, 8), 6.73 (2, s), 4.06 (2, q, 7), 2.12 (6, s), 1.36 (3, t, 7)
31	$\delta = 12.61$ (1, bs), 9.40 (1, bs), 9.06 (1, bs), 7.95 (1, bs), 7.69 (2, d, 8), 7.39 (2, d, 8), 6.90 (2, s), 5.20 (2, s), 2.16 (6, s)
32	$\delta = 12.65$ (1, bs), 9.42 (1, bs), 9.12 (1, bs), 7.97 (1, s), 7.71 (2, d, 8), 7.41 (2, d, 8), 7.10 (2, s), 2.88 (4, s), 2.15 (6, s)
33	$\delta = 13.21$ (1, bs), 9.88 (1, bs), 8.25 (1, s), 7.70 (1, d, 17), 7.60 (2, d, 9), 7.56 (2, s), 7.42 (2, d, 9), 6.49 (1, d, 17), 2.11 (6, s)
34	$\delta = 12.66$ (1, bs), 9.42 (1, bs), 9.28 (1, bs), 7.96 (1, s), 7.60 - 7.75 (3, m), 7.47 (2, s), 7.37 (2, d, 9), 6.46 (1, d, 17), 2.18 (6, s)
35	$\delta = 13.15$ (1, bs), 9.88 (1, bs), 8.24 (1, bs), 7.71 (1, d, 17), 7.62 (2, d, 8), 7.45 (2, d, 8), 7.23 (2, s), 6.66 (1, d, 17), 3.75 (6, s)

-87-

Co.No.	NMR results
36	$\delta = 13.10$ (1, bs), 9.84 (1, bs), 8.22 (1, s), 7.30-7.80 (8, m), 7.12 (1, bs), 6.65 (1, d, 17), 2.11 (6, s)
37	$\delta = 12.64$ (1, bs), 9.41 (1, bs), 9.23 (1, bs), 7.96 (1, s), 7.69 (2, d, 8), 7.51 (2, s), 7.46 (1, d, 15), 7.35 (2, d, 8), 7.20 (1, d, 15), 3.19 (3, s), 2.95 (3, s), 2.19 (6, s)
38	$\delta = 12.64$ (1, bs), 9.42 (1, bs), 9.23 (1, bs), 7.96 (1, bs), 7.67 (2, d, 8), 7.40-7.60 (3, m), 7.34 (2, d, 8), 7.26 (1, d, 15), 3.50-3.78 (8, m), 2.18 (6, s)
39	$\delta = 13.19$ (1, bs), 9.89 (1, bs), 8.25 (1, s), 7.45-7.65 (5, m), 7.39 (2, d, 9), 7.29 (1, d, 17), 3.60-3.80 (8, m), 2.11 (6, s)

D. Pharmacological example

The *in vitro* pharmacological activity of the present compounds was examined using one of the following rapid, sensitive and automated assay methods.

Method A

An HIV-1 transformed T4-cell line, MT-4, which was previously shown (Koyanagi et al., *Int. J. Cancer*, 36, 445-451, 1985) to be highly susceptible to and permissive for HIV infection, served as the target cell line. Inhibition of the HIV-induced cytopathic effect was used as the end point. The viability of both HIV- and mock-infected cells was assessed spectrophotometrically via the *in situ* reduction of 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT). The 50% cytotoxic concentration (CC₅₀ in M) was defined as the concentration of compound that reduced the absorbance of the mock-infected control sample by 50%. The percent protection achieved by the compound in HIV-infected cells was calculated by the following formula :

$$\frac{(\text{OD}_T)_{\text{HIV}} - (\text{OD}_C)_{\text{HIV}}}{(\text{OD}_C)_{\text{MOCK}} - (\text{OD}_C)_{\text{HIV}}} \quad \text{expressed in \%},$$

whereby (OD_T)_{HIV} is the optical density measured with a given concentration of the test compound in HIV-infected cells; (OD_C)_{HIV} is the optical density measured for the control untreated HIV-infected cells; (OD_C)_{MOCK} is the optical density measured for the control untreated mock-infected cells; all optical density values were determined at

-88-

540 nm. The dose achieving 50% protection according to the above formula was defined as the 50% effective concentration (EC₅₀ in M). The ratio of CC₅₀ to EC₅₀ was defined as the selectivity index (SI).

5 Method B

An HIV-1 transformed T4-cell line, MT-4, which was previously shown (Koyanagi et al., *Int. J. Cancer*, **36**, 445-451, 1985) to be highly susceptible to and permissive for HIV infection, served as the target cell line. In these cells, engineered with GFP (and an HIV-specific promotor), ongoing HIV-infection was measured fluorometrically.

10 Cytotoxicity is measured in the same cells, but engineered with GFP under a constitutional promotor. The infection (or inhibition thereof) of HIV infected cells and the fluorescence of mock-infected cells is assessed by the fluorescent GFP signal generated by the two above mentioned cell lines.

The 50% effective concentration (EC₅₀ in μ M) was defined as the concentration of
15 compound that reduced the fluorescence of HIV-infected cells by 50%. The 50% cytotoxic concentration (CC₅₀ in μ M) was defined as the concentration of compound that reduced fluorescence of the mock-infected cells by 50%. The ratio of CC₅₀ to EC₅₀ was defined as the selectivity index (SI).

20 The compounds of formula (I) were shown to inhibit HIV-1 effectively. Particular EC₅₀, CC₅₀ and SI values are listed in Table 6 hereinbelow.

Table 6 lists the pEC₅₀ (-logEC₅₀), pCC₅₀ (-logCC₅₀) and pSI (pCC₅₀-pEC₅₀) values for the compounds of formula (I). For example, a compound with a EC₅₀ value of 10⁻⁹M, i.e. pEC₅₀ = 9, and a CC₅₀ value of 10⁻⁵ M, i.e. pCC₅₀ = 5, has a SI of
25 10⁻⁵ M/10⁻⁹M = 10.000, i.e. a pSI of 5-9 = -4.

Table 6

Co. No.	pEC ₅₀ (M)	pCC ₅₀	pSI	Method
25	8.5	<4.5	<-4	A
23	8.2	<4.6	<-3.6	A
28	9.0	<4.6	<-4.4	A
24	8.4	4.3	-4.1	A
21	8.4	4.8	-3.6	B
4	8.4	5.0	-3.4	B

-89-

Co. No.	pEC50 (M)	pCC50	pSI	Method
20	8.4	5.2	-3.2	B
33	8.5	5.0	-3.5	B
13	8.2	5.2	-3.1	B
32	8.3	4.8	-3.5	B
26	9.3	5.0	-4.3	B
19	9.2	<4.6	<-4.6	B
3	8.5	5.9	-2.6	B
34	8.5	4.8	-3.7	B
35	8.4	4.9	-3.5	B
29	8.7	4.9	-3.8	B
37	8.1	5.4	-2.7	B
30	8.8	4.8	-4.0	B
1	8.4	<4.6	<-3.8	B
2	9.0	5.3	-3.7	B
5	8.5	5.3	-3.2	B
6	9.0	6.1	-2.9	B
31	9.3	<4.6	<-4.7	B
14	8.3	5.5	-2.8	B
54	8.7	5.5	3.2	B
55	8.6	5.6	3.0	B
45	9.2	<4.6	>4.6	B
46	9.4	<4.6	>4.8	B
72	8.5	<4.6	>3.9	B
64	9.3	<4.6	>4.7	B
53	8.7	6.1	2.6	B
42	8.0	<4.6	>3.4	B
62	8.6	<4.6	>4.0	B
43	8.0	<4.6	>3.4	B
52	9.3	5.7	3.6	B
48	9.2	<4.6	>4.6	B
50	9.2	5.8	3.4	B
63	8.4	5.1	3.3	B
49	9.0	5.5	3.5	B
44	8.0	4.7	3.3	B

-90-

Co. No.	pEC50 (M)	pCC50	pSI	Method
57	9.0	<4.6	>4.4	B
59	8.5	<4.6	>3.9	B
56	8.4	<4.6	>3.8	B
58	8.3	5.1	3.2	B
65	8.1	5.0	3.1	B
60	8.6	5.0	3.6	B
68	8.0	<4.6	>3.4	B
73	8.3	<4.6	>3.7	B